

# ESCMID<sup>†</sup> and ECMM<sup>‡</sup> joint clinical guidelines for the diagnosis and management of rare invasive yeast infections

M. C. Arendrup<sup>1</sup>, T. Boekhout<sup>2,3,4</sup>, M. Akova<sup>5</sup>, J. F. Meis<sup>6,7</sup>, O. A. Cornely<sup>8</sup>, O. Lortholary<sup>9,10</sup> and on behalf of the ESCMID EFISG study group and ECMM\*

1) Unit of Mycology, Department of Microbiological Surveillance and Research, Statens Serum Institut, Copenhagen, Denmark, 2) CBS Fungal Biodiversity Centre, Institute of the Royal Netherlands Academy of Arts and Sciences (KNAW), 3) Department of Internal Medicine and Infectious Diseases, University Medical Centre, Utrecht, the Netherlands, 4) Shanghai Key Laboratory of Molecular Medical Mycology, Changzheng Hospital, Second Military Medical University, Shanghai, China, 5) Section of Infectious Diseases, Hacettepe University School of Medicine, Ankara, Turkey, 6) Department of Medical Microbiology and Infectious Diseases, Canisius Wilhelmina Hospital, 7) Department of Medical Microbiology, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands, 8) 1st Department of Internal Medicine, Clinical Trials Centre Cologne, ZKS Köln, BMBF 01KN1106, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Cologne, Germany, 9) Centre National de Référence Mycoses Invasives et Antifongiques, Unité de Mycologie Moléculaire, Institut Pasteur, CNRS URA3012, and 10) Service des Maladies Infectieuses et Tropicales, Hôpital Necker-Enfants malades, APHP, Centre d'Infectiologie Necker-Pasteur, IHU Imagine, Université Paris Descartes, Paris, France

## Abstract

The mortality associated with invasive fungal infections remains high with that involving rare yeast pathogens other than *Candida* being no exception. This is in part due to the severe underlying conditions typically predisposing patients to these healthcare-related infections (most often severe neutropenia in patients with haematological malignancies), and in part due to the often challenging intrinsic susceptibility pattern of the pathogens that potentially leads to delayed appropriate antifungal treatment. A panel of experts of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Fungal Infection Study Group (EFISG) and the European Confederation of Medical Mycology (ECMM) undertook a data review and compiled guidelines for the diagnostic tests and procedures for detection and management of rare invasive yeast infections. The rare yeast pathogens were defined and limited to the following genera/species: *Cryptococcus adeliensis*, *Cryptococcus albidus*, *Cryptococcus curvatus*, *Cryptococcus flavescens*, *Cryptococcus laurentii* and *Cryptococcus uniguttulatus* (often published under the name *Filobasidium uniguttulatum*), *Malassezia furfur*, *Malassezia globosa*, *Malassezia pachydermatis* and *Malassezia restricta*, *Pseudozyma* spp., *Rhodotorula glutinis*, *Rhodotorula minuta* and *Rhodotorula mucilaginosa*, *Sporobolomyces* spp., *Trichosporon asahii*, *Trichosporon asteroides*, *Trichosporon dermatis*, *Trichosporon inkin*, *Trichosporon jirovecii*, *Trichosporon loubieri*, *Trichosporon mucoides* and *Trichosporon mycotoxinivorans* and ascomycetous ones: *Geotrichum candidum*, *Kodamaea ohmeri*, *Saccharomyces cerevisiae* (incl. *S. boulardii*) and *Saprochaete capitatae* (*Magnusiomyces* (*Blastoschizomyces*) *capitatus* formerly named *Trichosporon capitatum* or *Geotrichum* (*Dipodascus*) *capitatum*) and *Saprochaete clavata*. Recommendations about the microbiological investigation and detection of invasive infection were made and current knowledge on the most appropriate antifungal and supportive treatment was reviewed. In addition, remarks about antifungal susceptibility testing were made.

**Keywords:** Clinical guideline, *cryptococcus*, *Geotrichum*, *Kodamaea*, *Malassezia*, *Pseudozyma*, *Rhodotorula*, *Saccharomyces*, *Saprochaete*, *Sporobolomyces*, *Trichosporon*

**Article published online:** 18 September 2013

*Clin Microbiol Infect* 2014; **20** (Suppl. 3): 76–98

**Corresponding author:** M. C. Arendrup, Unit of Mycology, Statens Serum Institut (b. 43/317), Artillerivej 5, DK-2300 Copenhagen, Denmark

**E-mail:** maca@ssi.dk

and

Olivier Lortholary, Institut Pasteur, Centre National de Référence Mycoses Invasives et Antifongiques, Unité de Mycologie Moléculaire, CNRS URA3012, 75015 Paris, France

**E-mail:** olivier.lortholary@pasteur.fr

\*ESCMID/EFISG study group and ECMM members are listed in Appendix 1.

<sup>†</sup>European Society for Clinical Microbiology and Infectious Diseases. <sup>‡</sup>European Confederation of Medical Mycology.

## Introduction

In 2012 the first official European Society for Clinical Microbiology and Infectious Diseases (ESCMID) guideline on the diagnosis and treatment of a fungal infection was published [1–6]. Mucosal and invasive candidosis were covered and a comprehensive consensus guideline was developed with the participation of many experts from the ESCMID Fungal Infection Study Group (EFISG) representing many European countries. Before publication the recommendations were presented for discussion at a European Congress for Clinical Microbiology and Infectious Diseases (ECCMID) workshop and the subsequent manuscripts underwent peer-review before publication in the ESCMID journal, *Clinical Microbiology and Infection*.

Following the same rigorous procedure, EFISG continued the ESCMID guideline development process—this time in collaboration with the European Confederation of Medical Mycology (ECMM) and focusing on rare invasive fungal infections. The definition of such pathogens is somewhat pragmatic but yeasts other than *Candida*, mucorales, hyalohyphomycetous and dematiaceous fungi that are not common causes of invasive infections were included. This guideline presents the diagnostic and management guideline for ‘rare invasive yeast infections’ including several basidiomycetous yeasts: *Cryptococcus adeliensis*, *Cryptococcus albidus*, *Cryptococcus curvatus*, *Cryptococcus flavescens*, *Cryptococcus laurentii* and *Cryptococcus uniguttulatus* (often published under the name *Filobasidium uniguttulatum*), *Malassezia furfur*, *Malassezia globosa*, *Malassezia pachydermatis* and *Malassezia restricta*, *Pseudozyma* spp., *Rhodotorula glutinis*, *Rhodotorula minuta* and *Rhodotorula mucilaginoso*, *Sporobolomyces* spp., *Trichosporon asahii*, *Trichosporon asteroides*, *Trichosporon dermatis*, *Trichosporon inkin*, *Trichosporon jirovecii*, *Trichosporon loubieri*, *Trichosporon mucoides* and *Trichosporon mycotoxinivorans* and ascomycetous ones: *Geotrichum candidum*, *Kodamaea ohmeri*, *Saccharomyces cerevisiae* (incl. *S. boulardii*) and *Saprochaete capitatae* (*Magnusiomyces* (*Blastoschizomyces*) *capitatus* formerly named *Trichosporon capitatum* or *Geotrichum* (*Dipodascus*) *capitatum*) and *Saprochaete clavata*.

The selection of organisms has been based on the following criteria: (i) species were only included if they were documented as a cause of human invasive infections and (ii) rare *Candida* species were excluded because we anticipate that readers would probably refer to the ESCMID *Candida* guidelines rather than a rare invasive yeast infection guideline concerning species like *Candida palmioleophila* etc. although such species fulfil the term of being a rare cause of invasive infections. The species not considered being documented as

cause of invasive human disease included *Cryptococcus albidosimilis*, *Cryptococcus diffluens*, *Cryptococcus humicola* and *Cryptococcus uzbekistanensis*, *Trichosporon* spp. others than those mentioned above, *Blastobotrys proliferans*, *Millerozyma farinosa*, *Ogataea polymorpha* and *Guehomyces pullulans*. Finally, the guideline was limited to true yeasts and hence the unicellular algae *Prototheca wickerhamii* and *Prototheca zopfii* var. *zopfii* were excluded although we realize they have been misidentified as yeasts on occasion.

For the uncommon *Candida* species we refer the reader to the general recommendations regarding diagnosis and treatment as described in the ESCMID *Candida* guideline [1–6]. However, as some of these species are characterized by unique intrinsic susceptibility patterns, which are not specifically addressed in the *Candida* guidelines, a table summarizing this information has been elaborated and included here (Table 1, [7–19]). This table also includes names used in the anamorphic and teleomorphic states, despite this distinction recently being made superfluous [20].

General recommendations regarding collection, transport and storage of clinical specimens, direct examination, isolation and identification procedures, which are valid for all yeast-associated human infections, can be found in appropriate textbooks (e.g. Barnett et al. [21]) and are not mentioned here. Only specific features regarding genus identifications for the specific yeasts discussed herein were considered. The methods to evaluate the quality of evidence and to reach consensus recommendations were described previously [1]. Strength of recommendations’ quality of evidence was graded according to the criteria outlined in Table 2.

## Rare Invasive Yeast Infections

It is important to underscore that the fungal organisms covered in this guideline are not rare *per se*. A number of the ‘rare yeasts’ are encountered as frequent colonizers of human skin, mucosal surfaces, in food items or in the environment. In the normal host, infections are typically limited to various superficial infections like pityriasis versicolor, white piedra and occasionally onychomycosis, the management of which are dealt with in dermatology guidelines [22–25]. However, in the immunosuppressed or otherwise compromised host, invasive infections may occur, some being related to the presence of a central venous catheter (CVC) and a few reported as nosocomial clusters that require molecular approaches to be properly documented. As predicted from their low pathogenicity, invasive infections are still reported at low numbers in severely immunocompromised hosts (Table 3) [26–30]. For example, these organisms together constituted 1.1% of the

**TABLE 1.** Summary of rare *Candida* species that have been associated with human infection (by anamorphic and teleomorphic name). Remarks that may be relevant in the clinical context are included. References are kept to a minimum as these species are not the topic of this guideline

Anamorphic state	Teleomorphic state	Specific comment relevant in clinical context
<i>C. africana</i>	Not described	Closely related to <i>C. albicans</i> . Intrinsic susceptibility pattern as for this species. Probably less pathogenic than <i>C. albicans</i> and almost exclusively found in female genital tract specimens [7]
<i>C. auris</i>	Not described	Related to <i>C. haemulonii</i> . Fluconazole MICs higher than for <i>C. albicans</i> [8]
<i>C. bracarensis</i>	Not described	Closely related to <i>C. glabrata</i> . Susceptibility pattern as for <i>C. glabrata</i> (azole MICs elevated compared with <i>C. albicans</i> ) [9]
<i>C. ciferrii</i>	<i>Trichomonascus ciferrii</i>	Clinical significance uncertain. Inherent resistance to several antifungal compounds described [10]
<i>C. dubliniensis</i>	Not described	Closely related to <i>C. albicans</i> . Intrinsic susceptibility pattern as for this species. However, potential for acquired resistance to fluconazole appears to be greater than for <i>C. albicans</i> [11]
<i>C. fabianii</i>	<i>Cyberlindnera fabianii</i>	Clinical significance uncertain. Fluconazole MICs higher than for <i>C. albicans</i> [12]
<i>C. famata</i>	<i>Debaromyces hansenii</i>	This species has been reported as an infrequent cause of fungaemia. However, recent data questions if this species is actually human pathogenic (lack of growth at 37°C and no cases confirmed by sequencing [13])
<i>C. guilliermondii</i>	<i>Meyerozyma guilliermondii</i>	Closely related to <i>C. fermentati</i> and <i>C. palmiophila</i> . High echinocandin and azole MICs [13]
<i>C. haemulonii</i> (incl. <i>C. duobushaemulonii</i> )	Not described	Emerging evidence suggests that it may be a human pathogen related to superficial infections and central venous catheter-related fungaemia, particularly in Brazil, the Caribbean and Asian regions. Elevated azole and amphotericin MICs are reported. Related to <i>C. auris</i> [14]
<i>C. hellenica</i>	<i>Zygoascus meyeriae</i>	Fungaemia and respiratory infection has been reported. Decreased susceptibility to fluconazole, itraconazole, caspofungin, susceptible to voriconazole [15]
<i>C. inconspicua</i>	Not described	Closely related to <i>C. norvegensis</i> . Susceptibility pattern similar to <i>C. krusei</i> (intrinsically resistant to fluconazole) [16]
<i>C. intermedia</i>	Not described	Oropharyngeal colonizer, bloodstream infections, peritonitis. Susceptible to antifungal drugs except flucytosine [13]
<i>C. kefyr</i>	<i>Kluyveromyces marxianus</i>	No inherent resistance to antifungals described
<i>C. lipolytica</i>	<i>Yarrowia lipolytica</i>	Clinical significance uncertain. Fluconazole MICs higher than for <i>C. albicans</i> .
<i>C. lusitanae</i>	<i>Clavispora lusitanae</i>	Not a good target for amphotericin B even if MICs are in the susceptible range ( $\leq 1$ mg/L) [17]
<i>C. metapsilosis</i>	Not described	Closely related to <i>C. parapsilosis</i> . Susceptibility pattern similar to this species (high echinocandin MICs) [18]
<i>C. nivariensis</i>	Not described	Closely related to <i>C. glabrata</i> . Susceptibility pattern as for <i>C. glabrata</i> (decreased susceptibility to azoles) [9]
<i>C. norvegensis</i>	<i>Pichia norvegensis</i>	Closely related to <i>C. inconspicua</i> . Susceptibility pattern similar to <i>C. krusei</i> (intrinsically resistant to fluconazole) [16]
<i>C. orthopsilosis</i>	Not described	Closely related to <i>C. parapsilosis</i> . Susceptibility pattern similar to this species (high echinocandin MICs) [18]
<i>C. palmiophila</i>	Not described	Phenotypically related to <i>C. guilliermondii</i> . High azole MICs (but low echinocandin MICs in contrast to those for <i>C. guilliermondii</i> ) [13]
<i>C. pelliculosa</i>	<i>Wickerhamomyces anomala</i> (prev. <i>Pichia anomala</i> , <i>Hansenula anomala</i> )	Fluconazole MICs higher than for <i>C. albicans</i> [19]
<i>C. pulcherrima</i>	<i>Pichia kudriavzevii</i> (prev. <i>Metschnikowia pulcherrima</i> )	Clinical significance uncertain
<i>C. rugosa</i>	Not described	Fluconazole MICs higher than for <i>C. albicans</i> [19]
<i>C. subhashii</i>	Not described	Clinical significance uncertain
<i>C. viswanathii</i>	Not described	Clinical significance uncertain
<i>C. zeylanoides</i>	Not described	Clinical significance uncertain

**TABLE 2.** Strength of the EFISG and ECMM recommendation and quality of evidence

Strength of a recommendation (SoR)	
Grade A	ESCMID and ECMM strongly support a recommendation for use
Grade B	ESCMID and ECMM moderately support a recommendation for use
Grade C	ESCMID and ECMM marginally support a recommendation for use
Grade D	ESCMID and ECMM support a recommendation against use
Quality of evidence (QoE)	
Level I	Evidence from at least one properly designed randomized, controlled trial
Level II <sup>a</sup>	Evidence from at least one well-designed clinical trial, without randomization; from cohort or case–controlled analytical studies (preferably from more than one centre); from multiple time series; or from dramatic results of uncontrolled experiments
Level III	Evidence from opinions of respected authorities, based on clinical experience, descriptive case studies, or reports of expert committees
<sup>a</sup> Added index: ∙: Meta-analysis or systematic review of randomized controlled trials. ∙: Transferred evidence, that is, results from different patients' cohorts, or similar immune-status situation. ∙: Comparator group is a historical control. ∙: Uncontrolled trial. ∙: Published abstract (presented at an international symposium or meeting).	

almost 4000 fungaemia isolates in an 8-year national surveillance programme in Denmark [26,27]. In Paris, they represented 5.1% of the 3668 fungaemia isolates in a prospective surveillance programme from 2002 to 2012 (YEASTS Network, National Reference Centre for Invasive Mycoses and Antifungals, Paris, France (O. Lortholary, unpublished data), Table 3).

*Saccharomyces cerevisiae* is a biotechnologically highly important fungus with a broad use in the production of food and alcoholic beverages etc. Phylogenetically, the species is relatively closely related to *Candida glabrata* [31].

As most of these rare invasive yeast infections occur in the haematology and Intensive Care Unit (ICU) settings, clinicians should be aware that all of the responsible fungal species, presumably except *Saccharomyces* spp. and *Kodamaea ohmeri*, are regarded as intrinsically resistant to echinocandins. As a result of the rare nature of these pathogens, controlled prospective and comparative clinical trials are not feasible so solid data on treatment efficacy cannot be compiled. Moreover, clinical susceptibility breakpoints have not been established. Hence, management recommendations derive from clinical experience (cohort or case–controlled analytical studies, from multiple time series), pragmatic interpretations of susceptibility data and limited animal studies when available. Also because of their rare incidence, primary prophylaxis is not indicated unless specific local epidemiology suggests otherwise. As specific diagnostic surrogate markers have not been developed for these organisms (apart from the antigen test for *Cryptococcus* but mainly evaluated for *Cryptococcus*

**TABLE 3.** Summary of rare yeast isolates collected during the national surveillance programme in Denmark 2004–2011 and the surveillance programme in Paris hospitals, France October 2002–May 2012. Only unique isolates are included. For comparison and representing other parts of the world data from a US cancer centre, the Artemis study and a Brazilian study are included

	DK (national) [26,27]	Paris Hosp. <sup>a</sup>	US (cancer centre) [28]	Artemis study 1997–2007 [29]	Sao Paulo Brazil [30]
Fungaemia isolates (total)	3982	3668	3382	NA	1195
Rare yeasts other than <i>Candida</i>	44 (1.1%)	188 (5.1%)	94 (2.8%)	11,240	174 (14.5%)
<i>Cryptococcus neoformans</i>	13 (29.5%)	137 (72.8%)	NA	3,512 (31.2%)	79 (45.4%)
<i>Cryptococcus</i> spp.	1 (2.3%)	1 (0.5%)	NA	113 (1.0%)	NA
<i>Geotrichum</i> spp.	2 (4.5%)	19 (10.1%)	2 (5%)	NA	NA
<i>Rhodotorula</i> spp.	4 (9.1%)	5 (2.7%)	21 (51%)	462 (4.1%)	28 (16.1%)
<i>Saccharomyces</i> spp. <sup>b</sup>	22 (50.0%)	14 (7.4%)	8 (20%)	1,321 (11.8%)	NA
<i>Trichosporon</i> spp.	2 (4.5%)	11 (5.9%)	8 (20%)	1,196 (10.6%)	NA
<i>Malassezia</i> spp.	0	1	1 (2%)	NA	NA
<i>Pichia anomala</i>	0	NA	1 (2%)	28 (0.2%)	32 (18.4%)
<i>Saprochaete capitata</i>	0	NA	NA	109 (1.0%)	NA

NA, Not available.

<sup>a</sup>YEASTS Network, National Reference Centre for Invasive Mycoses and Antifungals, Paris, France (unpublished data).

<sup>b</sup>Taxonomically also a *Candida* species (*C. robusta*).

*neoformans*), blood culture remains an essential investigation for the detection of invasive infection and the general recommendations concerning volume and sampling frequency should be followed [32–34]. Emerging evidence has highlighted the additional yield obtained for candidaemia when a fungal blood culture bottle is included to specifically support the growth of fungi and at the same time avoid suppression by concomitant faster growing microorganisms [35–38]. It remains to be demonstrated if this applies to the detection of fungaemia due to the yeasts described herein. In the following sections the general characteristics for the different pathogens will be reviewed (in alphabetical order). Specific characteristics concerning the epidemiology and species identification are summarized in Tables 2 and 3. In most cases, identification to the species level requires the adoption of new tools including matrix-assisted laser desorption/ionization time of flight mass spectrometry (MALDI-TOF-MS) or DNA sequencing. In this context it is important to underline that the performance of these techniques depends on the quality of the databases employed. However, these approaches appear promising for most of these species and are expected to play an important role in the future (Table 4). Recommendations regarding use of blood culture and surrogate markers are summarized in Table 5. Finally, recommendations regarding appropriate first-line antifungal treatment options are summarized in Table 6.

## Cryptococcus

### Introduction

*Cryptococcus* is an anamorphic basidiomycetous yeast genus that comprises 70 species [39] but only two species *C. gattii* and *C. neoformans* are regularly causing infections. The diagnosis and

management of these have been described in detail elsewhere and are not included in the present guideline because they are not uncommon or rare infections [40–43].

Other *Cryptococcus* species (e.g. *C. albidus*, *C. curvatus*, *C. laurentii*) are prevalent worldwide and have been identified from various environmental sources including air, soil, water, pigeon droppings and food items such as cheese, milk, beans and wine [39]. The species are able to grow at 37°C and have been described as a cause of invasive human infections, with *C. albidus* and *C. laurentii* accounting for 80% of the non-*neoformans*/non-*gattii* cryptococcal infections [44]. *Cryptococcus uniguttulatus*, *C. adeliensis* and *C. flavescens* have also been implicated in cases of meningitis, albeit less frequently [45–48]. The clinical presentation is similar to that for *C. neoformans* but the cryptococcal antigen test is often negative and intrinsic susceptibility patterns are characterized by higher MICs for several agents [49]. Therefore, diagnosis and optimal management depend on a high index of suspicion and a skilled mycology laboratory service. In the literature, cultures were not always performed and species identifications were in many cases based solely on phenotypic methods. Therefore, reports on infections caused by the individual species should be interpreted with caution unless coming from reference laboratories.

### Risk factors and clinical presentation

*Cryptococcus adeliensis*, *C. albidus*, *C. curvatus*, *C. flavescens*, *C. laurentii* and *C. uniguttulatus* have been recovered from various clinical specimens [44–48,50]. The most common underlying risk factor is impaired cellular immunity, of which 16% was related to human immunodeficiency virus infection in a recent review of 44 previously published cases [44]. In addition, *C. laurentii* has been linked to the presence of invasive devices [44]. Clinical presentation involves bloodstream infection in 33–55% of the cases, neurological manifestations in 20–33%

**TABLE 4. Identification of the rare yeasts; laboratory characteristics and tools for identification to the species level**

Organism	Pseudo-hyphae	Hyphae	Blastoconidia	Anelloconidia	Chlamydoconidia	Cap-sule	Monopolar budding	Arthroconidia	Urease	Appearance on CHROMagar	Specific comments regarding morphology	Identification to the species level requires <sup>a</sup>
<i>Cryptococcus</i> spp. other than <i>C. neoformans</i> & <i>C. gattii</i>	-	-	+	-	-	+	-	-	+	No data	India ink may visualize the capsule in clinical specimens	ITS 1 + 2 (2nd option D1/D2 domain) MALDI-TOF promising
<i>Geotrichum</i>	-	+	-	-	+/(-)	-	-	+	-	Variable, probably species dependent	White colonies, arthroconidia	ITS 1 + 2 (2nd option D1/D2 domain) MALDI-TOF promising
<i>Kodamaea ohmeri</i>	+	-	+	-	-	-	-	-	-	Gradually pink to blue	Colour change on CHROMagar within 48 h	ITS 1 + 2 (2nd option D1/D2 domain) MALDI-TOF promising
<i>Malsessezia</i>	-	-	+	-	-	-	+	-	+	No growth except <i>M. pachydermatis</i>	Lipid dependent except <i>M. pachydermatis</i> white/cream colonies, monopolar budding	Biochemical tests or ITS 1 + 2 MALDI-TOF promising
<i>Pseudozyma</i>	-	+	+	-	-	-	-	-	+	<i>P. aphidis</i> forms rough green colonies. No data for the other species.	Fusiform spindle-shaped elongated blastoconidia	ITS 1 + 2 (2nd option D1/D2 domain)
<i>Rhodotorula</i>	-/(+)	-/(+)	+	-	-	(+)	-	-	+	No data	Red, orange-salmon coloured colonies	ITS 1 + 2 (2nd option D1/D2 domain) Biochemical tests or MALDI-TOF
<i>Saccharomyces</i>	-	-	+	-	-	-	-	-	-	Dark pink	Larger cells compared with <i>C. glabrata</i> , poor growth on blood agar	Biochemical tests or MALDI-TOF
<i>Saprocete capitata</i>	-	+	+/-	+	-	-	-	+	-	No data	Whitish, butyrous colonies, arthroconidia	ITS 1 + 2 (2nd option D1/D2 domain) MALDI-TOF promising
<i>Sporobolomyces</i>	-/(+)	-/(+)	+	-	-	-	-	-	+	No data	Red, orange-salmon coloured colonies; satellite colonies due to ballistoconidia	ITS 1 + 2 (2nd option D1/D2 domain)
<i>Trichosporon</i>	-	+	+	-	-	-	-	+	+	<i>T. asahii</i> ; dirty green (others variable)	Colony becoming dry, arthroconidia	ITS 1 + 2 (2nd option D1/D2 domain) IGS1 necessary for some species MALDI-TOF promising

<sup>a</sup>ITS 1 + 2: Sequencing of the Internal transcribed spacer 1 and 2 of the rDNA. 2nd option D1/D2 domain: Sequencing of the D1/D2 domain of the large subunit (LSU) ribosomal DNA 09FDNA recommended if species identification is not obtained by ITS 1 + 2 sequencing. IGS1: Sequencing of the intergenic spacer 1 region of the rDNA is necessary for correct species identification of some *Trichosporon* species. Biochemical tests: correct species identification can be obtained by biochemical tests such as the Vitek 2 and API ID32C. MALDI-TOF: matrix-assisted laser desorption/ionization time of flight mass spectrometry.

**TABLE 5.** Summary of and recommendations regarding use of blood culture and surrogate markers

Species	Surrogate markers	Strength of recommendation	Quality of evidence	Comment	References
All	Blood culture	A	II	Volume of blood is essential. Adults: 40–60 mL either one venepuncture or separate immediately after each other. Repeated if signs and symptoms of fungaemia persist.	[32–34]
	Blood culture fungal medium	B	II <sup>c</sup>	Several studies have documented better yield for <i>Candida</i> and BACTEC and BacT/ALERT BC systems if a mycosis medium is included. Not shown specifically for rare yeasts.	[26,35–38]
<i>Cryptococcus</i> other than <i>C. neoformans</i> and <i>C. gattii</i>	Cryptococcus antigen	C	III	Cerebrospinal fluid and serum. Sensitivity lower than for <i>C. neoformans</i> . (4/17 in one study). Negative result does not exclude cryptococcosis.	[44,45]
	Asp GM	D	III	Cross-reaction due to galactomannan in the fungal cell wall. However, sensitivity not examined.	[65]
<i>Geotrichum</i> <i>Kodamaea ohmeri</i> <i>Malassezia</i>	$\beta$ -D-glucan	D	II	$\beta$ -D-glucan is not part of the cryptococcal cell wall.	[63]
	No data on surrogate markers	No recommendation	No data		
	No data on surrogate markers	No recommendation	No data		
<i>Pseudozyma</i> <i>Rhodotorula</i>	Isolator 10 Lysis centrifugation or BC bottle suppl. with palmitic acid with prolonged incubation (2 weeks)	B	III	Positive microscopy and negative culture may be suggestive, due to the lipid dependence of most species. Subculture: Sabouraud overlaid with sterile olive oil, Dixon agar or other lipid-containing agar.	[107,108]
	No data on surrogate markers	No recommendation	No data		
<i>Saccharomyces</i>	$\beta$ -D-glucan	C	III	$\beta$ -D-glucan present in culture supernatant but in lower amount and but no clinical data.	[63]
	<i>Candida</i> mannan antigen	C	III	$\beta$ -D-glucan present in culture supernatant but no clinical data.	[63,180]
<i>Saprochaete capitata</i>	Asp GM	C	III	Single clinical case and antigen similarity	[179]
	Asp GM	C	III	Cross-reaction due to galactomannan in the fungal cell wall. However, sensitivity not examined.	[208,209]
<i>Trichosporon</i>	$\beta$ -D-glucan	C	III	$\beta$ -D-glucan present in culture supernatant but no clinical data.	[63]
	Cryptococcus antigen	B	II <sup>u</sup>	Cross-reaction with cryptococcal polysaccharide and galactomannan Ags. Dual positivity may be suggestive for <i>Trichosporon</i> infection. However, sensitivity not examined.	[57,58,246–249]
	Asp GM	C	III		
	$\beta$ -D-glucan	D	II	Low sensitivity	[238,250,251]

and pulmonary infection in 5–11% of the cases, but other body sites may also be involved including skin, eyes, peritoneum (secondary to peritoneal dialysis) and lymph nodes (5–10% each). *Cryptococcus laurentii* was associated with no mortality in contrast to *C. albidus* (28% mortality) [44].

### Diagnosis

Cryptococci are budding, encapsulated, round to oval yeast cells with a size from 3 to 8  $\mu$ m in fluids or tissues. The cells can be visualized by mixing the pellet (cerebrospinal fluid, pleural fluid or bronchoalveolar lavage) with India ink. The capsule surrounding the cell will appear transparent as a halo resembling an egg-white around the yolk. Tissue sections can be stained with mucicarmine or Alcian blue to highlight the capsule and so exclude other yeasts with similar morphology. Clinical specimens should be incubated on Sabouraud or other selective fungal agar at 30–35°C (for up to 4 weeks in case of prior antifungal treatment). Lysis centrifugation or culture of the buffy coat improves the yield of fungaemia detection [40]. Creamy mucoid colonies are obtained and the colour may darken with age. The genus is non-fermentative and identified by the presence of a capsule, urease production and assimilation pattern using commercial kits [39]. When cultured on bird seed agar most species do not form brown to black colonies (and can therefore be differentiated from *C. neoformans* and

*C. gattii*). There are few exceptions such as *Cryptotrichosporon anacardii* [51] and *Cryptococcus podzolicus* [52]. Species identification requires sequencing of internal transcribed spacer (ITS) 1 + 2 regions of rDNA and/or D1/D2 domains [39,47,48,53]; MALDI-TOF-MS is a promising tool for future identification of uncommon yeasts including *Cryptococcus* species; however, its performance will depend on the quality of the database (T. Boekhout, unpublished observations) [54–56].

Despite sharing several capsular antigens with *C. neoformans*, cryptococcal antigen detection is not a reliable diagnostic test for non-*neoformans*/non-*gattii* cryptococcal species with only 4/17 bloodstream cases being positive and at low titres [44,45]. Whether the poor sensitivity is attributable to differences in capsule structure (versus *C. neoformans*) or lower fungal burden remains to be clarified. False-positive results have been reported during invasive *Trichosporon* infections [57,58] and in the case of rheumatoid factor [59], *Capnocytophaga canimorsus* septicaemia [60], *Stomatococcus mucilaginosus* bacteraemia [61], soaps and disinfectants [62]. The amount of  $\beta$ -1-3-D-glucan in the cryptococcal cell wall is much lower than in for example *Candida albicans* and the  $\beta$ -1-3-D-glucan test has not been found useful in the diagnosis of cryptococcal disease, as exemplified by only two of 12 patients with cryptococcosis positive at a cut-off value of 80 pg/mL [63,64]. Therefore the test is not recommended in this context. Finally, cross-reaction with the

**TABLE 6. Targeted antifungal treatment of emerging invasive yeast infections. Note: intention is to cure manifest infection**

	Population/ manifestation	Antifungal	Strength of recommendation – quality of evidence	Comments	References
<i>Cryptococcus</i> other than <i>C. neoformans</i> and <i>C. gattii</i>	CNS and severe inf. Induction	Amphotericin <sup>a</sup> (±flucytosine <sup>b</sup> )	B-III	MICs of 5-FC, fluconazole and other azoles often elevated and particularly so for <i>C. albidus</i> , <i>C. laurentii</i> and <i>C. uniguttulatum</i> [29,30,33,54–59] If <i>in vitro</i> susceptible	[44–46,49,69–74]
	CNS and severe inf. Consolidation	Fluconazole ≥400 mg/day	C-III		
	Non-CNS, not severe inf.	Fluconazole ≥400 mg/day	C-III	MICs of 5-FC, fluconazole and other azoles often elevated and particularly so for <i>C. albidus</i> , <i>C. laurentii</i> and <i>C. uniguttulatum</i> [29,30,33,54–59]	
<i>Geotrichum</i> <i>candidum</i>	Non-CNS, not severe inf.	Amphotericin <sup>a</sup>	B-III	May be preferable to fluconazole for the less azole-susceptible species	
	Any	Echinocandins	D-II	Intrinsically resistant	
	Any	Amphotericin <sup>a</sup> (±flucytosine <sup>b</sup> )	B-III	Preferred agent is amphotericin B (w/wo 5-FC).	[77–80]
	Any	Fluconazole	D-III	No human data and high MIC values	[77,82]
<i>Kodamaea</i> <i>ohmeri</i>	Any	Voriconazole	NR	Low MIC values. One breakthrough failure case (while on micafungin), but voriconazole TDM levels not reported.	[77,78,81]
	Any	Echinocandins	D-II	<i>In vitro</i> resistant	[77]
	Any	Amphotericin <sup>a</sup>	B-III	Most (but limited) clinical experience w amphotericin B;	[83,85,86,92,95,96]
<i>Malassezia</i>	Any	Fluconazole	C-III	Elevated MICs for some isolates. Fluconazole successful in 5/6 paediatric cases, in 1/1 adult immunocompromised patient case (fluconazole followed by itraconazole) unsuccessful in one adult case of cellulitis [70,74,78].	[84,88,92]
	Any	Echinocandins	C-III	Two case reports showed successful outcome for one patient each on micafungin and caspofungin respectively (MICs higher than for <i>C. albicans</i> ) No data	[91,93]
	Any	Voriconazole	NR		
	Severe cases	Amphotericin <sup>a</sup>	B-III		[97,106]
	Non-severe cases	Fluconazole	B-III	<i>M. pachydermatis</i> may be less susceptible <i>in vitro</i> <i>In vitro</i> activity, no clinical data, exposure issues particularly in the paediatric population	[97,106]
	Any	Voriconazole	C-III	No reference susceptibility test for <i>Malassezia</i> ; modified susceptibility tests suggest intrinsic resistance	[97,106]
<i>Pseudozyma</i> spp.	Any	Flucytosine <sup>b</sup>	D-III	No reference susceptibility test for <i>Malassezia</i> ; modified susceptibility tests suggest intrinsic resistance	
	Any, fungaemia	Amphotericin <sup>a</sup>	A-II <sup>d</sup>	Low MICs of amphotericin B, case stories reporting success.	[121–124]
	Any, fungaemia	Fluconazole	D-II	5/6 clinical isolates had high MICs of fluconazole	[121–124]
	Any, fungaemia	Echinocandins	D-II	<i>In vitro</i> resistant	[123,124]
	Any, fungaemia	Flucytosine <sup>b</sup>	D-II	<i>In vitro</i> resistant (MIC >64 mg/L)	[121,123]
<i>Rhodotorula</i>	Any, fungaemia	Voriconazole	C-III	<i>In vitro</i> activity, no clinical data, exposure issues	[123,124]
	Any	Amphotericin <sup>a</sup> (±flucytosine <sup>b</sup> )	A-II	<i>In vitro</i> susceptible only to these two agents. Good response in the case reports available.	[131,153,154]
	Any	Azoles	D-II <sup>d</sup>	Breakthrough cases on fluconazole and echinocandins. <i>In vitro</i> resistant to all azoles and echinocandins.	[131,147,148,153,154]
<i>Saccharomyces</i>	Any	Echinocandins	D-II	Breakthrough cases on azoles and echinocandins. <i>In vitro</i> resistant	[147,153,154]
	Any	Amphotericin <sup>a</sup>	B-III	Most clinical experience; toxicity risk higher than for echinocandins	[178,182,183]
	Any	Echinocandins	C-III	Two successful cases in the literature (± neutropenic), no emergence of <i>S. cerevisiae</i> after intro of echinocandins as first line agents for candidaemia, two recent failure cases neutropenic (Arendrup MC unpublished data)	[27,182,183]
	Any	Amphotericin B <sup>a</sup> + Flucytosine <sup>b</sup>	B-III	Excellent <i>in vitro</i> susceptibility. May be used in severe cases or when penetration into an infected focus is challenging.	[70,170,181,184,185]
	Any	Fluconazole	D-III	Increased occurrence in patients exposed to fluconazole; high fluconazole MICs (similar to those for <i>C. glabrata</i> )	[26,178]
<i>Saprochaete</i> <i>capitata</i>	Any	Discontinuation of probiotics	A-III	Probiotic containing <i>S. bouldarii</i> has been documented to be the origin of systemic infections.	[175,176]
	Any	Amphotericin <sup>a</sup> (±flucytosine <sup>b</sup> )	B-III	Most (but limited) experience. <i>In vitro</i> susceptibility of amphotericin B in the intermediate range, failures in hepatosplenic infections reported on amphotericin B monotherapy.	[186,194,196]
	Any	Voriconazole	B-III	Less data available, but promising <i>in vitro</i> susceptibility	[186]
	Any	Echinocandins	D-II	<i>In vitro</i> resistant and a case report document breakthrough infections on an echinocandin.	[211,212]
	Any	Fluconazole	NR	<i>In vitro</i> resistant – but animal model data suggest activity	[211–213]
<i>Sporobolomyces</i>	Any	CSF or interferon- $\gamma$ in combination w antifungal treatment	C-III	Improved <i>in vitro</i> phagocytic activity has been observed	[194,195,206]
	Any	Amphotericin <sup>a</sup>	C-III	Possibly a treatment option but insufficient clinical data	[69,222,223,230]
	Any	Voriconazole	C-III	Possibly a treatment option but insufficient clinical data	[69,225,230]

Table 6 (Continued)

	Population/ manifestation	Antifungal	Strength of recommendation – quality of evidence	Comments	References
Trichosporon	Any	Echinocandins	D-II	<i>In vitro</i> resistant	[69,230]
	Any	Fluconazole	D-II	<i>In vitro</i> resistant	[69,230]
	Any	Voriconazole	B-III	Preferred therapy, but as data are scarce	[233,237,238,259,260]
	Any	Fluconazole	C-III	Some clinical evidence for usefulness, but also some isolates with higher MICs	[192,232,238]
	Any	Echinocandins	D-II	<i>In vitro</i> resistant	[237–239,256,262]
	Any	Amphotericin <sup>a</sup>	D-III	Low success rates on amphotericin B and <i>in vitro</i> resistance reported	[192,232,261,265–267,270]
	Any	Flucytosine <sup>b</sup>	D-III	<i>In vitro</i> resistant	

5-FC, flucytosine; CNS, central nervous system; inf., infection; TDM, therapeutic drug monitoring; w/wo, with/without.

<sup>a</sup>Amphotericin, wherever mentioned, includes amphotericin B deoxycholate and its lipid formulations. Lipid formulations are preferred due to lower toxicity and liposomal amphotericin B specifically whenever central nervous system penetration is warranted.

<sup>b</sup>Flucytosine, wherever mentioned, is a possible option particularly in cases where penetration issues (e.g. central nervous system infection) or severity, suggest combination therapy may improve outcome. Flucytosine should only be used in combination due to the risk of selection of resistance and therapeutic drug monitoring is highly recommended due to the narrow therapeutic index.

*Aspergillus* galactomannan antigen test (Platelia *Aspergillus*; BioRad, Marnes la Coquette, France) has been observed due to the presence of galactoxylomannan in the cryptococcal capsule [65]. A commercially available cryptococcal lateral flow antigen detection device (IMMY; Immuno-Mycologics, Inc., Norman, OK, USA), has recently been introduced as a point-of-care test for the early diagnosis of cryptococcosis. The test has proven effective both for serum, cerebrospinal fluid and urine samples in comparison with conventional techniques of antigen detection but the performance for the detection of infections due to cryptococci other than *C. neoformans* has not been evaluated [66–68].

#### Susceptibility testing and treatment

Cryptococci are intrinsically resistant to echinocandins. The optimal treatment for invasive infections due to other *Cryptococcus* species has not been established. Amphotericin B alone or in combination with flucytosine and fluconazole either alone or after induction therapy with amphotericin B has been used. Predictors for mortality were infection with *C. albidus* (rather than *C. laurentii*), age above 45 years and central nervous system involvement [44]. *In vitro* susceptibility testing suggests that *C. albidus*, *C. curvatus* and *C. laurentii* are susceptible to amphotericin B with MICs comparable to those for *C. neoformans* [49]. On the contrary, MICs of flucytosine, fluconazole and other azoles are in most studies elevated and particularly so for *C. albidus*, *C. laurentii* and *C. uniguttulatus*, suggesting that these organisms may be less susceptible [45,46,49,69–74]. Moreover, fluconazole resistance has been reported more frequently in patients with previous azole exposure (83% versus 50%) [44]. The clinical implication of these observations remains to be understood. However, initial induction therapy with amphotericin B until clear clinical improvement and careful evaluation of fluconazole susceptibility if step-down to this agent is considered appears to be a sound and safe strategy.

### *Geotrichum candidum*

*Geotrichum candidum* (*Galactomyces candidus*) is a filamentous ascomycetous yeast that forms arthroconidia and that has rarely been reported to be responsible for disseminated infections in the haemato-oncology setting [75–77]. It is closely related to *Saprochaete capitata* (*Magnusiomyces capitatus*) and *Saprochaete clavata* (see below). A recent literature review of 12 invasive cases reported since 1971 revealed that 8/12 patients had underlying malignancy [77]. It has been anecdotally reported as a cause of invasive skin infection [78]. The preferred agent for invasive infections is amphotericin B with or without concomitant flucytosine as most experience exists with this agent [77,79,80]. Voriconazole may be a promising agent, as suggested by good *in vitro* susceptibility, but only a single case is reported in the literature with an unsuccessful outcome of a breakthrough infection (while on micafungin) [77,78,81]. Neither echinocandins nor fluconazole can be recommended because of high MICs and no clinical support for efficacy [77,82].

### *Kodamaea ohmeri*

#### Introduction

*Kodamaea (Pichia) ohmeri* is a rarely occurring yeast that has recently been identified as a cause of fungaemia, endocarditis, cellulitis, funguria and peritonitis in neonates and children [83–87], and in both immunocompromised [88–91] and immunocompetent [92,93] adult patients. The anamorphic state is *Candida guilliermondii* var. *membranaefaciens*, and has been confused with *Candida guilliermondii* (for which the teleomorphic state is *Meyerozyma guilliermondii*). Recent literature invariably does not acknowledge the anamorphic state (and readers may therefore not recognize this species as a *Candida* sp.), so we have included *K. ohmeri* in this guideline. *Kodamaea ohmeri* may be

misidentified as *Candida tropicalis* [84], *Candida haemulonii* or *Candida parapsilosis* [94] by traditional methods and molecular identification is mandatory.

#### Risk factors/clinical presentation

There are <30 cases described in the world literature, which are mainly sporadic reports but also include two Asian series with several cases [84,94]. In one of these, the hands of a healthcare worker at a surgical ward were colonized with *K. ohmeri*, suggesting a nosocomial outbreak [84].

#### Diagnosis

Blood culture remains the cornerstone in the diagnosis of invasive infection. On solid media, *K. ohmeri* forms *Candida*-like colonies, which on CHROMagar (BioMérieux, Marcy l'Etoile, France) change colour from pink to blue within 48 h [88]. Although mis-classification as *Candida tropicalis* has been reported, correct species identification can be obtained by biochemical tests such as the Vitek 2 and API ID32C or sequencing of the rDNA loci [84,88].

#### Susceptibility testing and treatment

Susceptibility testing suggests that the susceptibility pattern mirrors that for *Candida glabrata* with azole MICs higher than those for *Candida albicans* (fluconazole: 2–64 mg/L; voriconazole: 0.03–8 mg/L, posaconazole 0.06–4 mg/L, respectively), but amphotericin B and echinocandin MICs in the range that would be interpreted as susceptible for *Candida albicans* (amphotericin B: 0.25–1 mg/L, caspofungin: 0.125–1 mg/L, micafungin: 0.03–0.06 mg/L, respectively) [84,94]. Most cases have been treated with liposomal amphotericin B (or amphotericin B deoxycholate) and with good response [83,85,86,92,95,96]. Fluconazole was successful in 5/6 paediatric cases, in 1/1 adult immunocompromised patient (fluconazole followed by itraconazole) and was associated with failure in one adult case of cellulitis [84,88,92]. Finally, caspofungin or micafungin treatment has been successful in one case each [91,93]. Hence, although there are insufficient data to support a firm treatment recommendation, amphotericin B appears to be an attractive first-line agent and echinocandins are possibly promising alternative candidates. Susceptibility testing is recommended not only to guide treatment but also to provide MIC–outcome relationships and hence data for future optimized treatment recommendations.

## Malassezia

### Introduction

*Malassezia* species are basidiomycetous yeasts and part of our normal skin microbiota. The genus includes 14 species of

which 13 are lipid dependent. These include *M. furfur*, *M. sympodialis*, *M. globosa*, *M. obtusa*, *M. restricta*, *M. slooffiae*, *M. dermatitis*, *M. japonica*, *M. nana*, *M. yamatoensis*, *M. equina*, *M. caprae* and *M. cuniculi*. *Malassezia pachydermatis*, however, is able to grow on routine media without the addition of oil or other sources of lipid. *Malassezia sympodialis*, *M. globosa*, *M. slooffiae* and *M. restricta* are the most frequently found species responsible for colonization of humans.

#### Risk factors/clinical presentation

*Malassezia* species may cause various skin manifestations including pityriasis versicolor, seborrhoeic dermatitis, dandruff, atopic eczema and folliculitis and less commonly onychomycosis [97]. These skin manifestations are common and are typically diagnosed and managed in the primary healthcare sector or by dermatologists [97]. For their management, we would refer the reader to local and international guidelines. However, skin colonization and infection may be a source for transmission to vulnerable patient groups susceptible to invasive infections [97]. Both the non-lipid-dependent species *M. pachydermatis* and the lipid-dependent *Malassezia* spp. (almost all reported as caused by *M. furfur*) have been reported to cause systemic infections [97–99]. As an example, *M. pachydermatis*, which is known to cause external otitis in dogs, has been isolated from the hands of dog owners, including healthcare providers such as nurses, and associated with clusters of infections in neonates [98,100–102]. Apart from contact with a potential carrier, other risk factors for *M. pachydermatis* include increased median neonatal acute physiology score and more than 9 days of arterial catheterization [98]. Notably, *M. pachydermatis* can persist for a long time on surfaces of incubators, which may serve as a source of infection for neonates [103].

Systemic infections due to lipid-dependent *Malassezia* species mainly occur in the following host groups: (i) infants on lipid-containing parenteral nutrition and (ii) children and adults with various forms of immunosuppression and underlying diseases [97,104]. Risk factors for fungaemia related to non-lipid total parenteral nutrition are the following: chronic ambulatory peritoneal dialysis (due to lipids leaking from the gastrointestinal tract), haematological malignancy, cancer and Crohn's disease and most cases arise in patients with a CVC [97,105,106].

#### Diagnosis

The diagnosis of invasive *Malassezia* infections is challenging because of the lipid-dependent nature of most species. Hence, special media, such as modified Dixon or modified Leeming and Notham agar, or the use of Sabouraud agar with the addition of a few drops of sterile olive oil is required. The

performance of the various modern automated blood culture systems for the detection of *Malassezia* species has not been systematically studied, however, the use of the Isolator 10 system (with subculture of lipid-containing agar) or supplementation of the blood culture flask with palmitic acid has been shown to be required for the detection of lipid-dependent species in earlier blood culture systems [107,108]. Inoculated blood culture media should be incubated for up to 2 weeks. Colonies on solid agar are cream to beige. Yeast cells are round or oval with thick walls with a size varying from 2 to 8 µm. Cell division is via monopolar budding with buds that may be nearly as broad as the mother cell in many cases, leading to a flask-like appearance of the cells.

*Malassezia* species may be distinguished phenotypically using morphology and a series of biochemical tests as well as using molecular tools [109–111] and by MALDI-TOF-MS (T. Boekhout, unpublished observation). Full protocols for phenotypic identification are provided by Guého-Kellermann et al. [112]. For clinical management at the level of the individual patient, species identification is less important, although it is obviously needed for epidemiological surveillance and outbreak investigation.

#### Susceptibility testing and treatment

Susceptibility testing of *Malassezia* has not been standardized because growth is not supported on the standard RPMI growth medium recommended for yeast and mould testing by CLSI and EUCAST. Significant variation and broad range MICs have been reported in various publications depending on the medium used, which may result in random/erroneous susceptibility classification [113]. Some reports suggest that *in vitro* fluconazole resistance may be encountered more often in *M. pachydermatis* [114,115]. The clinical implication of this finding remains unclear. So far susceptibility testing for guiding treatment cannot be recommended. As for the other rare invasive yeast infections, larger patient series are lacking, and hence evidence-based treatment recommendations cannot be made. The key factors in the management of invasive *Malassezia* infection are removal of the CVC, discontinuation of the parenteral lipid and institution of systemic antifungal treatment. Most experience is with fluconazole and amphotericin B, which are the preferred agents [106,116,117]. In general MICs are lower for voriconazole than for fluconazole for most species, however, so is the voriconazole exposure, particularly in the paediatric population, which is a reason to strongly consider therapeutic drug monitoring if voriconazole is prescribed [118–120]. Additionally, voriconazole is associated with more side-effects and drug–drug interactions and is not licensed for neonates or children <2 years old. Future studies are

warranted to elucidate which antifungal treatment is optimal. *In vitro* resistance to flucytosine and echinocandins appears to be a consistent finding and therefore these drug classes are not recommended [101,116].

## *Pseudozyma*

#### Introduction

*Pseudozyma* species are basidiomycetous plant pathogens, which belong to the Ustilaginales. The genus today contains at least 20 species and was not recognized as a human pathogen until 2003 when three *Pseudozyma* species—*Pseudozyma antarctica*, *Pseudozyma parantarctica* and *Pseudozyma thailandica*—were isolated from the blood of three Thai patients [121]. Fungaemia due to *Pseudozyma aphidis* was subsequently reported from the USA in 2008 [122] and recently in a neonate from India [123].

#### Risk factors/clinical presentation

So far infections due to *Pseudozyma* have been reported from Asia (Korea, China, India, Thailand), Brazil and the USA [123]. Risk factors associated with invasive *Pseudozyma* infections are similar to those of non-*albicans* *Candida* spp., i.e. extremes of age, cancer chemotherapy, neutropenia (<3000 cells/µL), presence of a CVC and severe thrombocytopenia [121–124]. Invasive *Pseudozyma* spp. infection most often presents with fungaemia. Moreover, deep-seated focal infections including brain abscess [125] and pleural cavity have been reported [124]. Finally, a single case of a mycetoma due to co-infection with *Nocardia* after traumatic inoculation in an Asian farmer has been described [126].

#### Diagnosis

The main stay in the diagnosis of invasive *Pseudozyma* spp. infection is blood culturing and when, focal infection is suspected, culture of relevant tissue samples. On Sabouraud agar *Pseudozyma* species form rapidly expanding moist, tan-yellow and wrinkled yeast colonies [122,124] and on CHROMagar *Candida* medium *P. aphidis* forms rough green colonies after 48–72 h of incubation at 37°C (no data for the other species). Microscopic examination shows fusiform spindle-shaped elongated blastoconidia and the presence of hyphae. Germ tube test and chlamydoconidia formation are negative. Isolates show positive test for diazonium blue B and hydrolyse urea. Growth is inhibited on 0.1% cycloheximide-containing medium. API ID 32C and VITEK2 compact generally give non-conclusive profile(s). Amplification and sequencing of the ITS and/or D1/D2 domain is necessary for a proper identification [121–123,125,126].

### Susceptibility testing and treatment

Low MIC values of amphotericin B, posaconazole (0.03 mg/L), voriconazole (0.06 mg/L) and isavuconazole (0.25 mg/L) have been reported. Susceptibility of itraconazole was variable and 5/6 isolates had MICs of fluconazole in the range of 4 to greater than 64 mg/L. The echinocandins (>4 mg/L) and flucytosine (>64 mg/L) are not active against *Pseudozyma* [123]. Although the available data are too limited to provide firm treatment recommendations, first-line options may be amphotericin B or voriconazole whereas echinocandins, fluconazole and flucytosine should be avoided.

## Rhodotorula

### Introduction

Clinically relevant red yeasts belong to two genera: *Rhodotorula* and *Sporobolomyces* (see below). *Rhodotorula* species are common environmental basidiomycetous yeasts, which can be found in soil, ocean and lake water, fruit juice and milk, and on shower curtains and toothbrushes [30,127]. Today, the genus contains 46 species [128] of which three have been described as rare human pathogens: *R. mucilaginosa* (also known as *R. rubra*), *R. glutinis* and *R. minuta* [30,129,130]. The association of *R. mucilaginosa* with humans is well documented and this yeast has been isolated from skin, sputum and digestive tract samples including faeces, forming part of the normal human microbiota. This species accounts for the majority of the infections (74–79%) followed by *R. glutinis* (7.7%) [130,131]. In a significant proportion of the reported cases the species identification is not available (17%); furthermore, species identification is not reliable by methods normally available in the routine microbiology laboratory, and identification using sequence analysis of the DID2 domains of the large subunit ribosomal DNA (rDNA) and the ITS 1 + 2 regions of the rDNA is needed [30,130].

### Risk factors/clinical presentation

Opportunistic fungal infections due to *Rhodotorula* have emerged after the first case was reported in 1985, and the most common predisposing factor appears to be the presence of a CVC and underlying haematological disease [28,127,130–135]. However, *Rhodotorula* fungaemia, peritonitis, endocarditis or meningitis have also been reported in other vulnerable patient groups, including patients with AIDS, extensive burns, continuous ambulatory peritoneal dialysis, cirrhosis, those who have undergone intra-abdominal surgery, intravenous drug abusers and critically ill ICU patients [130,136–146]. Notably, a significant number of cases are breakthrough infections during fluconazole or echinocandin treatment [131,134,135,147,148]. Infections appear to be less common in Nordic countries

compared with the warmer regions [26,28,149]. Moreover, *Rhodotorula* has also been found on the hands of healthcare workers in Egypt [150], and it has a high affinity to adhere to plastic surfaces and can form biofilms [28,151]. Hence, medical equipment including flexible endoscopes, various utensils and furniture in the patient's room can easily become colonized. These observations suggest geographical differences in the epidemiology and possibly a potential role of differences in hygiene procedures [30,148,150].

### Diagnosis

The mainstay in the diagnosis of invasive *Rhodotorula* spp. infection is blood culture as 79% of the systemic infections presents as fungaemia [130]. Of note, isolates of *Rhodotorula* have been found to cross react with the *Candida glabrata*/*Candida krusei* probe in the commercially available fluorescence *in situ* hybridization test for presumptive species identification of positive blood cultures, which might lead to inappropriate echinocandin treatment [152]. A CVC is often involved and therefore the catheter tip should be cultured when the CVC is removed to capture cases where blood cultures are falsely negative [127]. *Rhodotorula* isolates are easily recognizable in the laboratory by their distinctive orange to salmon-coloured mucoid colonies. Cells of *Rhodotorula* spp. are subglobose, oval to elongate, with or without small capsules, may sometimes form rudimentary hyphae but not ballistoconidia, produce urease but fail to ferment carbohydrates.

Antigen detection has not been reported in clinical cases of systemic *Rhodotorula* infections.  $\beta$ -1-3-D-glucan has been detected in the supernatant from three isolates of *R. mucilaginosa* (syn. *R. rubra*) at an average concentration of two-thirds of that of *Candida* spp. [63]. Whether the  $\beta$ -1-3-D-glucan test would be useful as a surrogate marker for invasive *Rhodotorula* infection remains to be investigated.

### Susceptibility testing and treatment

Susceptibility testing has yielded amphotericin B MICs of <1 mg/L and flucytosine MICs of <0.5 mg/L, but fluconazole MICs of >32 mg/L and voriconazole, itraconazole and posaconazole median MICs of 2 mg/L, which is above the breakpoints for not only *Candida albicans* but also *Aspergillus fumigatus* [153–157]. Hence, *Rhodotorula* species are regarded as intrinsically resistant to azoles and echinocandins, but susceptible to amphotericin B and flucytosine [153,158,159]. This is further supported by the finding that many cases have been breakthrough infections during fluconazole or echinocandin treatment [131,147,148]. Consequently, the preferred treatment of choice is with any kind of amphotericin B preparation. With such treatment, an overall mortality was reported as 13.8% in the haematological setting and ranges

from 0% in patients with non-Hodgkin's lymphoma to 21% in patients with acute leukaemia [30,127,131,160]. If present, withdrawal of CVC is strongly recommended [130–133,135,147] and sometimes reported to be efficacious without any accompanying antifungal treatment [28,161]. Furthermore, cases of chronic ambulatory peritoneal dialysis-associated *Rhodotorula* peritonitis have been diagnosed and successfully treated (5/6 patients cured) with catheter removal (5/6 patients) and systemic or intraperitoneal antifungal therapy (amphotericin B in 5/6 patients or ketoconazole 1/6 patients, which is today regarded as an obsolete compound for oral administration, in one case) [130,138,162,163]. There are anecdotal reports indicating improvement of fungal infection under fluconazole and miconazole treatment despite *in vitro* resistance and in patients without antifungal treatment [130,164]. However, since there are no confirmatory trials for treatment of *Rhodotorula* infections, such practice should be avoided, especially in seriously ill patients.

## Saccharomyces

### Introduction

*Saccharomyces cerevisiae*, also known as baker's or brewer's yeast, is a low pathogenic ascomycetous yeast. *Saccharomyces boulardii*, a genetically similar subtype [165,166], is used as a probiotic for prevention and treatment of various sorts of diarrhoea and recurrent *Clostridium difficile*-associated diarrhoea [167–169] and should be avoided in immunocompromised hosts. The anamorphic state of *S. cerevisiae* is sometimes referred to as *Candida robusta*. As the species is closely related to *Candida glabrata* phylogenetically it is not surprising that the clinical and microbiological characteristics are similar to this species.

### Risk factors/clinical presentation

*Saccharomyces cerevisiae* may be found as a harmless and transient digestive commensal and colonizer of mucosal surfaces of normal individuals. It can, however, also be involved in mucosal infections like vaginitis, particularly in fluconazole-exposed women with recurrent vulvo-vaginal candidiasis, and in bloodstream infections, again particularly in fluconazole-exposed patients [35,170]. Cases of fungaemia and disseminated infection have been described in vulnerable patients after treatment with the *S. boulardii* probiotic compound [171–176], and also as nosocomial infection in a patient that shared a room with a patient receiving probiotic, suggesting transfer via contaminated hands of the nursing staff [177]. In a recent review of 92 cases of invasive *Saccharomyces* infections, *S. boulardii* accounted for half of these, was less often associated with an underlying immunocompromised

condition and more often associated with a favourable outcome [178]. Use of probiotics in debilitated patients, in ICU, in neutropenic patients, in preterm newborns or in patients with central lines should be carefully considered.

### Diagnosis

Invasive infection is most often diagnosed by microscopy and culture. Yeast cells are round to oval and larger than *Candida glabrata* cells. Ascospores are occasionally seen and short pseudohyphae may be formed but are not typical; urease activity is absent. *Candida* mannan antigen positivity has been anecdotally reported in patients with fungaemia but it remains unclear if this test can be used for diagnosis [179]. Similarly,  $\beta$ -1-3-D-glucan has been found in culture supernatant from *S. cerevisiae* at quantities of approximately 85% compared with that from *Candida* cultures and this test has been reported positive in case reports of *Saccharomyces* bloodstream infection and on culture supernatants from *S. cerevisiae* cultures, but the diagnostic performance has not been systematically studied [63,180].

### Susceptibility testing and treatment

The *in vitro* susceptibility pattern is similar to that of *Candida glabrata*, with elevated azole MICs, but echinocandin MICs only a few dilutions higher than for *Candida albicans* and low amphotericin B and flucytosine MICs [70,170,181]. Most clinical experience exists with fluconazole and amphotericin B, for which favourable outcome was observed for 60% and 77.7%, respectively [178]. The clinical experience with the echinocandin class of drugs is limited. Two cases have been reported in the literature and were successfully treated [182,183]. However, two recent failures, one of which with autopsy documented multi-organ dissemination, have also been observed (M. C. Arendrup, unpublished observations). Finally, amphotericin B with or without flucytosine has been used in severe or recurrent cases [184,185] but the role of this combination remains to be established. In addition to the systemic antifungal therapy, it is strongly recommended that probiotics containing *S. boulardii* are discontinued and indwelling foreign bodies are removed, when possible, because this organism, like many other yeasts, is capable of forming biofilms [176].

## Saprochaete

### Introduction

*Saprochaete capitata* (Teleomorph: *Magnusiomyces capitatus*, previously named *Geotrichum capitatum*, *Trichosporon capitatum* or *Blastoschizomyces capitatus*) is a non-fermentative, non-encapsulated, urease-negative ascomycetous yeast. It is found in environmental sources such as wood and soil, in

animals (including bovine mastitis and poultry faeces) and has been found in dishwashers [133,186–188]. In addition, *S. capitata* is part of the normal microbiota of human skin and is frequently isolated from sputum and the digestive tract of healthy people [186].

#### Risk factors/clinical presentation

*Saprochaete capitata* is a rare, but emerging yeast mostly responsible for often lethal fungaemia in patients with profound neutropenia in the haematology setting [186,189–191]. This patient category represents up to 92% of reported cases and 75% of them have been reported from Italy, Spain and France [192]. Mortality associated with disseminated infections has been estimated to be 57% in the haematology population [192,193]. Of note, *Geotrichum* spp. represented 5% (2/41) of non-*Candida*, non-*Cryptococcus* fungaemia cases among haematological patients in a tertiary cancer centre in Houston, suggesting that this species is still a rare invasive pathogen [28]. In the haemato-oncology setting, 60–80% of patients present with deep organ involvement [193]. Indeed, it has been recognized as a cause of skin lesions similar to those observed during disseminated candidiasis, hepatosplenic abscesses [194,195], pancreatic infections [196], brain abscesses [195], funguria [197], acute renal failure due to fungal occlusion of glomeruli [198] and osteomyelitis, mostly with vertebral involvement [199–201]. Central venous catheters have been recognized as a potential portal of entry [193].

A common hospital source has been advocated for several clusters [186,202] and more recently, it has been confirmed by sequencing that *S. capitata* in milk vacuum flasks was the origin of an outbreak in four patients in Barcelona [203]. No subsequent cases occurred when the identified contaminated source was withdrawn [203].

Outside the oncology context, *S. capitata* has been responsible for prosthetic valve endocarditis [204], pneumonia [205], fungaemia due to contaminated intravenous fluid [206] and meningitis [207].

#### Diagnosis

Most cases of *S. capitata* fungaemia have been diagnosed by means of blood culture. On solid media, *S. capitata* colonies are white to cream and isolates produce true hyphae, pseudohyphae, blastoconidia, arthroconidia and annelloconidia. *In vitro* and *in vivo* studies have revealed that *Magnusiomyces capitatus* antigens may cross-react with the *Aspergillus* galactomannan assay [208,209].  $\beta$ -1-3-D-glucan can be detected *in vitro* in culture supernatant at amounts of 88% compared with that for *Candida* spp. [63], but there is no experience of glucan detection during invasive human cases. The species can also be identified by MALDI-TOF-MS (T. Boekhout, unpublished observation).

#### Susceptibility testing and treatment

*In vitro* susceptibility data suggest that *S. capitata* is susceptible to flucytosine (MIC values 0.25–0.5 mg/L), itraconazole, voriconazole and posaconazole (MIC ranges: 0.12–0.50, 0.25–0.5 and 0.03–0.25 mg/L, respectively), but not to fluconazole (MIC between 16 and 32 mg/L) [193,202,210]. The MICs for amphotericin B ranged between 0.5 and 2.0 [186]. *Saprochaete capitata* can be considered intrinsically resistant to echinocandins and *S. capitata* was the cause of at least five reported episodes of breakthrough infection in neutropenic patients receiving echinocandins [211,212]. Finally, various combination regimens were not superior to high-dose fluconazole in an experimental animal model of *Blastoschizomyces capitatus* infection (MICs of 8 and 16 mg/L) [213]. The clinical implication of the conflicting observations regarding *in vitro* and *in vivo* activity of fluconazole remains to be understood.

There are not enough clinical data to assess the optimal treatment for *S. capitata* in haematology patients. However, based on *in vitro* data and the limited clinical data available, any amphotericin B formulation with or without flucytosine can be recommended [186,196]. Failure despite high-dose liposomal amphotericin B (7 mg/kg) has been reported in the context of hepatosplenic infection and neutropenic sepsis [191,194]. Voriconazole exhibits a promising activity *in vitro* [186] and some authors have suggested the use of voriconazole and amphotericin B combination therapy [193,214]. Of note, in the above-mentioned animal model, high-dose fluconazole was more efficacious than amphotericin B, flucytosine or voriconazole monotherapy [213]. Although echinocandin MIC values are elevated, isolated case reports have suggested the potential interest of the combination of caspofungin and voriconazole [215,216]. The role of echinocandins as part of combination therapy for *S. capitata* infections remains to be clarified.

Early removal of the catheter is an important complementary treatment as it has been the likely source of the infection in some cases [186,189] and as removal was shown as a prognostic indicator for success in one study [193]. Other adjuvant therapies to improve the phagocytic activity such as colony-stimulating factors, granulocyte transfusions and interferon- $\gamma$  have been combined with antifungal drugs with some success [194,195,217].

*Saprochaete clavata* (*Geotrichum clavatum*), which is closely related to *S. capitata*, has only very infrequently been described as involved in invasive human infection. However, 33 cases of *S. clavata* invasive infections were recently reported in France from January 2009 to August 2012, of which 17 cases were diagnosed within 2 months as part of an outbreak in haematology wards in 2012 (Vaux, ECCMID 2013,

O 505). It remains to be determined if the low number of reported cases is the result of difficult species identification and confusion with related species or if it is indeed a very uncommon infection outside isolated outbreaks.

## Sporobolomyces

### Introduction

*Sporobolomyces* species are usually red-to-orange pigmented basidiomycetous yeasts, that, next to regular budding cells, form ballistoconidia. Phylogenetically, they are closely related to *Rhodotorula* species. The 53 species occur widely in nature, especially on leaf surfaces, but also in soil, fruits etc. and *Sporobolomyces* is detected in indoor air, particularly in the summer season [218–221].

### Risk factors/clinical presentation

*Sporobolomyces* have been reported as the cause of sporadic invasive bloodstream infections particularly in AIDS patients [222,223]. A single case of meningitis due to *Sporobolomyces roseus* has been described in an immunocompetent cocaine abuser presenting with a 1-week history of severe headache and neck stiffness but no cerebrospinal fluid pleocytosis [224]. The cerebrospinal fluid was culture negative but the yeast was detected by molecular assays in two separate cerebrospinal fluid specimens. Amphotericin B has been efficacious in fungaemic and meningitis cases [222,224]. A single case of endogenous endophthalmitis due to *Sporobolomyces salmonicolor* in a patient with a history of pelvic inflammatory disease 2 years earlier has been reported [225]. The patient recovered following intravitreal amphotericin B (5 µg) and systemic voriconazole 200 mg twice daily. Finally, dermatitis due to *Sporobolomyces holsaticus* and allergic respiratory disease linked to *Sporobolomyces* exposure have been reported [226–228].

### Diagnosis

Colonies of *Sporobolomyces* species are usually red or orange in colour and are similar to those of *Rhodotorula* species. They differ from the latter by the formation of ballistoconidia that are actively discharged, which usually lead to the formation of many small satellite colonies. The optimal growth temperature is 25–30°C. Some isolates may fail to grow well at 35–37°C [218]. Species identification requires sequence analysis of the D1/D2 domains and the ITS1 + 2 regions of the rDNA. The performance of indirect tests such as  $\beta$ -1-3-D-glucan has not been investigated. Due to the ubiquitous presence of *Sporobolomyces* contamination, transient colonization or pseudoinfection should be considered [223,224,229].

### Susceptibility testing and treatment

*In vitro* susceptibility testing has been investigated for a limited number of *Sporobolomyces salmonicolor* isolates and suggests that this species is intrinsically resistant to fluconazole and echinocandins (MIC ranges 8–256 and  $\geq$ 128 mg/L of fluconazole and micafungin, respectively), but susceptible to voriconazole and terbinafine (0.03–2 and 0.06–0.12 mg/L, respectively) [69]. The MIC range of amphotericin B and itraconazole was 0.5–8 and 0.03–4 mg/L, respectively, suggesting variable susceptibility [69,230]. Although the available data are too limited to provide firm treatment recommendations, first-line options may be amphotericin B or voriconazole whereas echinocandins and fluconazole should be avoided [69,222,223,225,230]. Susceptibility testing is recommended.

## Trichosporon

### Introduction

*Trichosporon* species are urease-positive, non-encapsulated basidiomycetous yeasts with no known sexual state. They are widely distributed in the environment and regularly found on normal skin, particularly in the peri-genital areas, and occasionally as part of the normal gastrointestinal or upper respiratory microflora [133,231]. The most characteristic morphological feature is the formation of cylinder-shaped arthroconidia in addition to pseudohyphae, septate hyphae and blastoconidia. The genus has undergone a major taxonomic reclassification and today 37 species are described that belong to at least five phylogenetic clades [231]. However, only 16 species have been associated with human infection [231]. The vast majority of cases is caused by *Trichosporon asahii* (74%), followed by *T. dermatis* (12%) [232]. An obvious challenge in this respect is that a significant part of the existing literature does not provide a correct or unique species identification and hence species-specific data are limited.

### Risk factors/clinical presentation

In humans, *Trichosporon* has been associated with white piedra and hypersensitivity pneumonia particularly in hot and humid climates (16–30%). In the immunocompromised host, invasive infections such as fungaemia, endocarditis, peritonitis and meningitis have also been reported. Most common risk factors for invasive infections are underlying malignant haematological disease with long-term neutropenia [192,232,233] or with neutrophil dysfunction such as chronic granulomatous disease [234]. Finally, a recent report suggests that *T. mycotoxinivorans* may be an emerging pulmonary pathogen in patients with cystic fibrosis [235]. Males more often contract the infection and predisposing factors are the presence of a CVC, ICU stay,

peritoneal dialysis, steroid use and cytotoxic chemotherapy [192,231,232,236]. Previous other fungal systemic infection is not uncommon and breakthrough cases in patients receiving fluconazole or echinocandin have been described [237–239]. The infection most commonly presents as fungaemia (75%), in approximately 50% of the cases associated with metastatic skin lesions [232,233]. Renal involvement may occur and be associated with haematuria and funguria [232]. In immunocompromised patients, *Trichosporon* is increasingly seen at some centres among invasive yeast infections other than *Candida* and *Cryptococcus* and associated with a mortality rate of up to 80% [236,240,241], though 55% has been reported in a more recent report from Taiwan [232]. Of note, rare clinical cases have been reported in neonates and in intravenous drug abusers [242–244].

### Diagnosis

The cornerstone in the diagnosis of invasive infection is microscopy (for the detection of fungal disease) and culture. On solid media, colonies are white, but on CHROMagar *T. asahii* forms characteristic dirty green colonies. *Trichosporon* species share antigens with *Cryptococcus* and *Aspergillus* and a number of reports have demonstrated cross-reaction for the cryptococcal antigen and/or galactomannan antigen kits [57,58,245–249]. Therefore, dual positivity in these tests may be an indicator of invasive trichosporonosis; however, the sensitivity and specificity of this approach has not been defined [57,231,245]. On the other hand the  $\beta$ -1-3-D-glucan tests have been associated with a low diagnostic sensitivity for trichosporonosis [245,250,251]. Molecular tests, including direct detection on blood or formalin-fixed paraffin-embedded tissue samples, are being developed but are not yet standardized [252–255]. Reliable species identification requires molecular identification with sequencing of the ITS 1 + 2 ( $\pm$ D1/D2 domain) or even the intergenic spacer region (IGS1) of the rDNA [256–258]. MALDI-TOF-MS appears to be a promising identification tool (with an extensive database) [259].

### Susceptibility testing and treatment

Emerging experience suggests that azoles are the primary drug class for the treatment of invasive trichosporonosis [192,232,233,237,238,260,261]. Several of the species are resistant *in vitro* to amphotericin B with MICs  $\geq 2$  mg/L, including *T. asahii* [256,257,262,263]. *Trichosporon* species are resistant to flucytosine (MICs 4–128 mg/L) and to the echinocandins (MICs  $>16$  mg/L) [239,263,264]. In patients with systemic trichosporonosis and underlying haematological disease poor response rates (i.e. between 16% and 24%) on amphotericin B have been reported and therefore this agent is not recommended for invasive infections [192,231,265–267].

Triazoles on the other hand have been found to be superior to other antifungal drug classes in prophylaxis and treatment [232,238,268]. Most published experience concerns the use of fluconazole. However, variations in susceptibility *in vitro* may suggest that not all species and isolates are equally susceptible to this agent [257]. Voriconazole is the preferred agent because it displays good *in vitro* activity against most *Trichosporon* species and isolates and has been associated with good *in vivo* outcome in most cases of clinical and animal studies [232,233,256,261,263,269–271]. In addition to triazole treatment, resolution of myelosuppression and removal of vascular catheters are other confounders related with increased survival [28,192,231,237,238].

### Conclusion

Rare yeasts other than *Candida* and *Cryptococcus neoformans/gattii* are commonly found in the environment and as skin or mucosal colonizers in humans. Some of them may be considered as true emergent opportunistic pathogens in Europe, although the number of reported episodes remains low because of their low pathogenicity even in patients with severely compromised immunity, particularly those with haematological malignancies and a CVC.

Several of these rare yeasts possess intrinsic or variable resistance to antifungals including echinocandins (to which only *Saccharomyces* spp. and *K. ohmeri* are presumably susceptible) and even polyenes (*Trichosporon* spp.) or azoles (*Rhodotorula* spp.; some *M. capitatus* isolates). This may explain the occurrence of breakthrough infections during empiric antifungal therapy in neutropenic patients. Amphotericin B is among the recommended first-line treatment options for these infections except for *Trichosporon*. Combination with flucytosine may be considered particularly in severe cases or cases where drug penetration may be suboptimal provided the infecting organism is susceptible. In agreement with the recommendations provided in the *Candida* guideline amphotericin B lipid formulations may be considered preferable to conventional deoxycholate amphotericin B for toxicity reasons and liposomal amphotericin B particularly for infections involving the CNS [3,5,6,272,273].

Genus identification is mandatory for clinical management and should be performed and provided in a timely manner. Species identification of these rare yeasts, however, remains difficult and often requires reference expertise and adoption of modern techniques including molecular analysis or MALDI-TOF-MS (with an extensive database). Whether or not the precise species identification (outside cryptococcosis) may

have an impact on the individual clinical management remains to be documented; however, its use and value for epidemiological surveillance and outbreak investigation has been documented beyond doubt.

## Acknowledgements

The work behind this guideline has been supported by grants from ESCMID and ECMM. The guideline draft has been presented in part at the ECCMID conference 2013 in Berlin. TB and JFM have been supported by a Qatar National Research Fund grant NPRP 5-298-3-086.

## Transparency Declarations

MCA has received grant support from Astellas Pharma, Gilead Sciences, Merck Sharp and Dohme (MSD), Pfizer and Schering Plough. She has been a consultant or at the advisory board for Gilead Sciences, MSD, Pfizer, Pcovery and Schering Plough. She has been paid for talks on behalf of Gilead Sciences, MSD, Pfizer, Astellas Pharma and Schering Plough.

TB has no conflict of interest to declare.

MA has received grants and speaker's fees from Pfizer, Gilead and Merck Sharp and Dohme.

JFM is a consultant for Astellas, Basilea, MSD and Merck, and received grants or speaker's fees from Basilea, MSD, Merck, Schering Plough and Astellas.

OL is a consultant for Novartis, Fab Pharma and Gilead Sciences, and received grants or speaker's fees from MSD, Roche, Astellas, Gilead Sciences and Pfizer.

JG has received grant support from Astellas Pharma, Basilea Pharmaceutica and Fondo de investigación sanitaria (FIS). He has been paid for talks on behalf of Gilead Sciences, Pfizer, Astellas, Hikma Pharma, United Medical and MSD.

AJU has received research grants from MSD (Schering-Plough), and is/was an advisor or received lecture honorarium from Astellas, Aicuris, Basilea, Gilead, MSD and Pfizer.

SAA has received research grants from Pfizer and lecture honoraria from Merck and Pfizer.

PM has been advisor and received speaker's fees from Astellas, Pfizer, Gilead, Merck Sharp and Dohme and Novartis.

FL has received research grant from Gilead and speaker's fees from Gilead and MSD.

OAC is supported by the German Federal Ministry of Research and Education (BMBF 01KN1106), has received research grants from 3M, Actelion, Astellas, Basilea, Bayer, Celgene, Cubist, F2G, Genzyme, Gilead, GSK, Merck/MSD, Miltenyi, Optimizer, Pfizer, Quintiles and Viropharma, is a

consultant to 3M, Astellas, Basilea, Cubist, F2G, Gilead, GSK, Merck/MSD, Optimizer, Pfizer and Sanofi Pasteur, and received lecture honoraria from Astellas, Gilead, Merck/MSD and Pfizer.

LP has received honoraria from Gilead Sciences, Schering-Plough, Astellas Pharma, Merck and Pfizer Pharmaceuticals, he has been speaker for Gilead Sciences, Schering-Plough, Merck, Pfizer Pharmaceuticals and Astellas Pharma.

JM has received research grants and lecture honoraria from Gilead, MSD (Schering-Plough) and Pfizer.

## Appendix I: ESCMID/EFISG Study Group and ECMM Members Contributing to this Guideline

Sevtap Arikan-Akdagli, Department of Medical Microbiology, Mycology Laboratory, Hacettepe University Medical School, Ankara, Turkey; Manuel Cuenca-Estrella, Servicio de Micología, Centro Nacional de Microbiología, Instituto de Salud Carlos III, Madrid, Spain; Eric Dannaoui, Unité de Parasitologie-Mycologie, Service de Microbiologie, Hôpital Européen G. Pompidou, Université Paris Descartes, APHP, Paris, France; Anne D. van Diepeningen, CBS Fungal Biodiversity Centre, Institute of the Royal Netherlands Academy of Arts and Sciences(KNAW), Utrecht, The Netherlands; Andreas H. Groll, Infectious Disease Research Program, Centre for Bone Marrow Transplantation and Department of Paediatric Haematology/Oncology, University Children's Hospital, Münster, Germany; Josep Guarro, Mycology Unit, Medical School & IISPV, Rovira i Virgili University, Reus, Spain; Jesús Guinea, Servicio de Microbiología Clínica y Enfermedades Infecciosas Clinical Microbiology and Infectious Diseases, Hospital General Universitario Gregorio Marañón, Madrid, Spain; William Hope, Department of Molecular and Clinical Pharmacology, The University of Liverpool, Liverpool, UK; Michaela Lackner and Cornelia Lass-Flörl, Division of Hygiene and Medical Microbiology, Innsbruck Medical University, Innsbruck, Austria; Katrien Lagrou, KU Leuven, Department of Microbiology and Immunology, Clinical Department of Laboratory Medicine, Leuven, Belgium; Fanny Lanternier, Unité de Mycologie Moléculaire, Institut Pasteur, Centre National de Référence Mycoses Invasives et Antifongiques, CNRS URA3012 and Université Paris Descartes, Service des Maladies Infectieuses et Tropicales, Hôpital Necker-Enfants Malades, APHP, Centre d'Infectiologie Necker-Pasteur, IHU Imagine, Paris, France; Joseph Meletiadis, Clinical Microbiology Laboratory, Attikon University Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece; Patricia Munoz, Departamento de Medicina, Clinical Microbiology and Infectious Diseases, Hospital General Universitario Gregorio Marañón, Universidad Complutense, Madrid, Spain; Livio Pagano, Istituto di Ematologia, Università

cattolica S. Cuore, Roma, Italy; Malcolm D. Richardson Mycology Reference Centre, National Aspergillus Centre, University Hospital of South Manchester, and Manchester Academic Health Science Centre, University of Manchester, Manchester, UK; Emmanuel Roilides, Infectious Diseases Unit, 3rd Department of Pediatrics, Hipokraton Hospital, Aristotle University, Thessaloniki, Greece, Anna Maria Tortorano, Department of Biomedical Sciences for Health, Università degli Studi di Milano, Milano, Italy; Andrew J. Ullmann, Division of Infectious Diseases, Department of Internal Medicine II, Julius-Maximilians-University, Würzburg, Germany.

## References

- Ullmann AJ, Cornely OA, Donnelly JP *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: developing European guidelines in clinical microbiology and infectious diseases. *Clin Microbiol Infect* 2012; 18: 1–8.
- Cuenca-Estrella M, Verweij PE, Arendrup MC *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: diagnostic procedures. *Clin Microbiol Infect* 2012; 18: 9–18.
- Cornely OA, Bassetti M, Calandra T *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: non-neutropenic adult patients. *Clin Microbiol Infect* 2012; 18: 19–37.
- Hope WW, Castagnola E, Groll AH *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: prevention and management of invasive infections in neonates and children caused by *Candida* spp. *Clin Microbiol Infect* 2012; 18: 38–52.
- Ullmann AJ, Akova M, Herbrecht R *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: adults with haematological malignancies and after haematopoietic stem cell transplantation (HCT). *Clin Microbiol Infect* 2012; 18: 53–67.
- Lortholary O, Petrikos G, Akova M *et al.* ESCMID\* guideline for the diagnosis and management of *Candida* diseases 2012: patients with HIV infection or AIDS. *Clin Microbiol Infect* 2012; 18: 68–77.
- Borman AM, Szekely A, Linton CJ, Palmer MD, Brown P, Johnson EM. Epidemiology, antifungal susceptibility, and pathogenicity of *Candida africana* isolates from the United Kingdom. *J Clin Microbiol* 2013; 51: 967–972.
- Lee WG, Shin JH, Uh Y *et al.* First three reported cases of nosocomial fungemia caused by *Candida auris*. *J Clin Microbiol* 2011; 49: 3139–3142.
- Mirhendi H, Bruun B, Schonheyder HC *et al.* Differentiation of *Candida glabrata*, *C. nivariensis* and *C. braccarensis* based on fragment length polymorphism of ITS1 and ITS2 and restriction fragment length polymorphism of ITS and D1/D2 regions in rDNA. *Eur J Clin Microbiol Infect Dis* 2011; 30: 1409–1416.
- Agin H, Ayhan Y, Devrim I, Gulfidan G, Tulumoglu S, Kayserili E. Fluconazole-, amphotericin-B-, caspofungin-, and anidulafungin-resistant *Candida ciferrii*: an unknown cause of systemic mycosis in a child. *Mycopathologia* 2011; 172: 237–239.
- Sullivan DJ, Moran GP, Pinjon E *et al.* Comparison of the epidemiology, drug resistance mechanisms, and virulence of *Candida dubliniensis* and *Candida albicans*. *FEMS Yeast Res* 2004; 4: 369–376.
- Bhally HS, Jain S, Shields C, Halsey N, Cristofalo E, Merz WG. Infection in a neonate caused by *Pichia fabianii*: importance of molecular identification. *Med Mycol* 2006; 44: 185–187.
- Jensen RH, Arendrup MC. *Candida palmioleophila*: characterisation of a previously overlooked pathogen and its unique susceptibility profile in comparison with five related species. *J Clin Microbiol* 2011; 49: 549–556.
- Cendejas-Bueno E, Kolecka A, Alastruey-Izquierdo A *et al.* Reclassification of the *Candida haemulonii* complex as *Candida haemulonii* (*C. haemulonii* group I), *C. duobushaemulonii* sp. nov. (*C. haemulonii* group II), and *C. haemulonii* var. *vulnera* var. nov.: three multiresistant human pathogenic yeasts. *J Clin Microbiol* 2012; 50: 3641–3651.
- Brandt ME, Kauffman CA, Pappas PG *et al.* Fungemia caused by *Zygoascus hellenicus* in an allogeneic stem cell transplant recipient. *J Clin Microbiol* 2004; 42: 3363–3365.
- Sugita T, Takeo K, Ohkusu M *et al.* Fluconazole-resistant pathogens *Candida inconspicua* and *C. norvegensis*: DNA sequence diversity of the rRNA intergenic spacer region, antifungal drug susceptibility, and extracellular enzyme production. *Microbiol Immunol* 2004; 48: 761–766.
- Atkinson BJ, Lewis RE, Kontoyiannis DP. *Candida lusitanae* fungemia in cancer patients: risk factors for amphotericin B failure and outcome. *Med Mycol* 2008; 46: 541–546.
- Mirhendi H, Bruun B, Schonheyder HC *et al.* Molecular screening for *Candida orthopsilosis* and *Candida metapsilosis* among Danish *Candida parapsilosis* group blood culture isolates: proposal of a new RFLP profile for differentiation. *J Med Microbiol* 2010; 59: 414–420.
- Pfaller MA, Diekema DJ, Messer SA, Boyken L, Hollis RJ, Jones RN. *In vitro* activities of voriconazole, posaconazole, and four licensed systemic antifungal agents against *Candida* species infrequently isolated from blood. *J Clin Microbiol* 2003; 41: 78–83.
- Taylor JW. One Fungus = One Name: DNA and fungal nomenclature twenty years after PCR. *IMA Fungus* 2011; 2: 113–120.
- Barnett JA, Payne RW, Yarrow D. *Yeasts: characteristics and identification*, 3rd edn. Cambridge: Cambridge University Press, 2007.
- Gupta AK, Ryder JE, Nicol K, Cooper EA. Superficial fungal infections: an update on pityriasis versicolor, seborrheic dermatitis, tinea capitis, and onychomycosis. *Clin Dermatol* 2003; 21: 417–425.
- Hu SW, Bigby M. Pityriasis versicolor: a systematic review of interventions. *Arch Dermatol* 2010; 146: 1132–1140.
- Khandpur S, Reddy BS. Itraconazole therapy for white piedra affecting scalp hair. *J Am Acad Dermatol* 2002; 47: 415–418.
- Bonifaz A, Gomez-Daza F, Paredes V, Ponce RM. Tinea versicolor, tinea nigra, white piedra, and black piedra. *Clin Dermatol* 2010; 28: 140–145.
- Arendrup MC, Bruun B, Christensen JJ *et al.* National surveillance of fungemia in Denmark 2004–2009. *J Clin Microbiol* 2011; 49: 325–334.
- Arendrup MC, Dzajic E, Jensen RH *et al.* Epidemiological changes with potential implication for antifungal prescription recommendations for fungaemia: data from a nationwide fungaemia surveillance programme. *Clin Microbiol Infect* 2013; 19: E343–E353.
- Chitasombat MN, Kofteridis DP, Jiang Y, Tarrand J, Lewis RE, Kontoyiannis DP. Rare opportunistic (non-*Candida*, non-*Cryptococcus*) yeast bloodstream infections in patients with cancer. *J Infect* 2012; 64: 68–75.
- Pfaller MA, Diekema DJ, Gibbs DL *et al.* Results from the ARTEMIS DISK global antifungal surveillance study, 1997 to 2007: 10.5-year analysis of susceptibilities of noncandidal yeast species to fluconazole and voriconazole determined by CLSI standardized disk diffusion testing. *J Clin Microbiol* 2009; 47: 117–123.
- Duboc De Almeida GM, Figueiredo Costa S, Melhem M *et al.* *Rhodotorula* spp. isolated from blood cultures: clinical and microbiological aspects. *Med Mycol* 2008; 46: 547–556.
- Kurtzman CP. Discussion of teleomorphic and anamorphic ascomycetous yeasts and yeast-like taxa. In: Kurtzman CP, Fell JW, Boekhout T, eds. *The yeasts, a taxonomic study*, 5th edn. Amsterdam: Elsevier, 2011; 293–307.
- Baron EJ, Weinstein MP, Dunne WM, Yagupsky P, Welsh DF, Wilson DM. *Cumitech 1C, blood cultures IV*. Washington, DC: ASM Press, 2005.

33. Arendrup MC, Bille J, Danaoui E, Ruhnke M, Heussel CP, Kibbler C. ECIL-3 classical diagnostic procedures for the diagnosis of invasive fungal diseases in patients with leukaemia. *Bone Marrow Transplant* 2012; 47: 1030–1045.
34. Arendrup M, Jensen IP, Justesen T. Diagnosing bacteremia at a Danish hospital using one early large blood volume for culture. *Scand J Infect Dis* 1996; 28: 609–614.
35. Arendrup MC, Sulim S, Holm A et al. Diagnostic issues, clinical characteristics, and outcomes for patients with fungemia. *J Clin Microbiol* 2011; 49: 3300–3308.
36. Ericson EL, Klingspor L, Ullberg M, Ozenci V. Clinical comparison of the Bactec Mycosis IC/F, BacT/Alert FA, and BacT/Alert FN blood culture vials for the detection of candidemia. *Diagn Microbiol Infect Dis* 2012; 73: 153–156.
37. Horvath LL, George BJ, Murray CK, Harrison LS, Hospenthal DR. Direct comparison of the BACTEC 9240 and BacT/ALERT 3D automated blood culture systems for candida growth detection. *J Clin Microbiol* 2004; 42: 115–118.
38. Cateau E, Cognee AS, Tran TC et al. Impact of yeast–bacteria coinfection on the detection of *Candida* sp. in an automated blood culture system. *Diagn Microbiol Infect Dis* 2012; 72: 328–331.
39. Fonseca A, Fell JW, Boekhout T. *Cryptococcus Vuillemin* (1901). In: Kurtzman CP, Fell JW, Boekhout T, eds. *The yeasts, a taxonomic study*, 5th edn. Amsterdam: Elsevier, 2011; 1616–1737.
40. Dromer F, Lortholary O. Cryptococcosis. In: Magill A, Ryan E, Solomon T, Hill D, eds. *Hunter's tropical medicine and emerging infectious disease*, 9th edn. Amsterdam: Elsevier, 2012; 1214.
41. Perfect JR, Dismukes WE, Dromer F et al. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2010; 50: 291–322.
42. Singh N, Dromer F, Perfect JR, Lortholary O. Cryptococcosis in solid organ transplant recipients: current state of the science. *Clin Infect Dis* 2008; 47: 1321–1327.
43. Jarvis JN, Dromer F, Harrison TS, Lortholary O. Managing cryptococcosis in the immunocompromised host. *Curr Opin Infect Dis* 2008; 21: 596–603.
44. Khawcharoenporn T, Apisarnthanarak A, Mundy LM. Non-neoformans cryptococcal infections: a systematic review. *Infection* 2007; 35: 51–58.
45. McCurdy LH, Morrow JD. Infections due to non-neoformans cryptococcal species. *Compr Ther* 2003; 29: 95–101.
46. Pan W, Liao W, Hagen F et al. Meningitis caused by *Filobasidium uniguttulatum*: case report and overview of the literature. *Mycoses* 2012; 55: 105–109.
47. Tintelnot K, Losert H. Isolation of *Cryptococcus adeliensis* from clinical samples and the environment in Germany. *J Clin Microbiol* 2005; 43: 1007.
48. Rimek D, Haase G, Luck A, Casper J, Podbielski A. First report of a case of meningitis caused by *Cryptococcus adeliensis* in a patient with acute myeloid leukemia. *J Clin Microbiol* 2004; 42: 481–483.
49. Bernal-Martinez L, Gomez-Lopez A, Castelli MV et al. Susceptibility profile of clinical isolates of non-*Cryptococcus neoformans*/non-*Cryptococcus gattii* *Cryptococcus* species and literature review. *Med Mycol* 2010; 48: 90–96.
50. Burnik C, Altintas ND, Ozkaya G et al. Acute respiratory distress syndrome due to *Cryptococcus albidus* pneumonia: case report and review of the literature. *Med Mycol* 2007; 45: 469–473.
51. Okoli I, Oyeka CA, Kwon-Chung KJ et al. *Cryptotrichosporon anacardii* gen. nov., sp. nov., a new trichosporonoid capsulate basidiomycetous yeast from Nigeria that is able to form melanin on niger seed agar. *FEMS Yeast Res* 2007; 7: 339–350.
52. Petter R, Kang B, Boekhout T, Davis B, Kwon-Chung K. A survey of heterobasidiomycetous yeasts for the presence of the genes homologous to virulence factors of *Filobasidiella neoformans*, CNLAC1 and CAP59. *Microbiology* 2001; 147: 2029–2036.
53. Bovers M, Diaz MR, Hagen F et al. Identification of genotypically diverse *Cryptococcus neoformans* and *Cryptococcus gattii* isolates by Luminex xMAP technology. *J Clin Microbiol* 2007; 45: 1874–1883.
54. Firacative C, Trilles L, Meyer W. MALDI-TOF MS enables the rapid identification of the major molecular types within the *Cryptococcus neoformans*/*C. gattii* species complex. *PLoS ONE* 2012; 7: e37566.
55. McTaggart LR, Lei E, Richardson SE, Hoang L, Fothergill A, Zhang SX. Rapid identification of *Cryptococcus neoformans* and *Cryptococcus gattii* by matrix-assisted laser desorption/ionization-time of flight mass spectrometry. *J Clin Microbiol* 2011; 49: 3050–3053.
56. Posteraro B, Vella A, Cogliati M et al. Matrix-assisted laser desorption/ionization–time of flight mass spectrometry-based method for discrimination between molecular types of *Cryptococcus neoformans* and *Cryptococcus gattii*. *J Clin Microbiol* 2012; 50: 2472–2476.
57. Fekkar A, Brun S, D'Ussel M et al. Serum cross-reactivity with *Aspergillus* Galactomannan and Cryptococcal antigen during fatal disseminated *Trichosporon dermatis* infection. *Clin Infect Dis* 2009; 49: 1457–1458.
58. Lyman CA, Devi SJ, Nathanson J, Frasch CE, Pizzo PA, Walsh TJ. Detection and quantitation of the glucuronoxylomannan-like polysaccharide antigen from clinical and nonclinical isolates of *Trichosporon beigelii* and implications for pathogenicity. *J Clin Microbiol* 1995; 33: 126–130.
59. Yeo SF, Wong B. Current status of nonculture methods for diagnosis of invasive fungal infections. *Clin Microbiol Rev* 2002; 15: 465–484.
60. Westerink MA, Amsterdam D, Petell RJ, Stram MN, Apicella MA. Septicemia due to DF-2. Cause of a false-positive cryptococcal latex agglutination result. *Am J Med* 1987; 83: 155–158.
61. Chanock SJ, Toltzis P, Wilson C. Cross-reactivity between *Stomatococcus mucilaginosus* and latex agglutination for cryptococcal antigen. *Lancet* 1993; 342: 1119–1120.
62. Blevins LB, Fenn J, Segal H, Newcomb-Gayman P, Carroll KC. False-positive cryptococcal antigen latex agglutination caused by disinfectants and soaps. *J Clin Microbiol* 1995; 33: 1674–1675.
63. Odabasi Z, Paetznick VL, Rodriguez JR, Chen E, McGinnis MR, Ostrosky-Zeichner L. Differences in  $\beta$ -glucan levels in culture supernatants of a variety of fungi. *Med Mycol* 2006; 44: 267–272.
64. Ostrosky-Zeichner L, Alexander BD, Kett DH et al. Multicenter clinical evaluation of the (1 $\rightarrow$ 3)  $\beta$ -D-glucan assay as an aid to diagnosis of fungal infections in humans. *Clin Infect Dis* 2005; 41: 654–659.
65. Dalle F, Charles PE, Blanc K et al. *Cryptococcus neoformans* galactoxylomannan contains an epitope(s) that is cross-reactive with *Aspergillus* galactomannan. *J Clin Microbiol* 2005; 43: 2929–2931.
66. McMullan BJ, Halliday C, Sorrell TC et al. Clinical utility of the cryptococcal antigen lateral flow assay in a diagnostic mycology laboratory. *PLoS ONE* 2012; 7: e49541.
67. Hansen J, Slechta ES, Gates-Hollingsworth MA et al. Large scale evaluation of the Immuno-Mycologics Inc. (IMMY) lateral flow and enzyme-linked immunoassays for the detection of cryptococcal antigen in serum and cerebrospinal fluid. *Clin Vaccine Immunol* 2013; 20: 52–55.
68. Jarvis JN, Percival A, Bauman S et al. Evaluation of a novel point-of-care cryptococcal antigen test on serum, plasma, and urine from patients with HIV-associated cryptococcal meningitis. *Clin Infect Dis* 2011; 53: 1019–1023.
69. Serena C, Pastor FJ, Ortoneda M, Capilla J, Noland N, Guarro J. *In vitro* antifungal susceptibilities of uncommon basidiomycetous yeasts. *Antimicrob Agents Chemother* 2004; 48: 2724–2726.
70. Quindos G, Ruesga MT, Martin-Mazuelos E et al. *In-vitro* activity of 5-fluorocytosine against 1,021 Spanish clinical isolates of *Candida* and other medically important yeasts. *Rev Iberoam Micol* 2004; 21: 63–69.
71. Shimokawa O, Niimi M, Kikuchi K, Saito M, Kajiwara H, Yoshida S. Relationship between MIC and minimum sterol 14 $\alpha$ -demethylation-inhibitory concentration as a factor in evaluating activities of

- azoles against various fungal species. *J Clin Microbiol* 2005; 43: 5547–5549.
72. Pedroso RS, Ferreira JC, Candido RC. *In vitro* susceptibility to antifungal agents of environmental *Cryptococcus* spp. isolated in the city of Ribeirao Preto, Sao Paulo, Brazil. *Mem Inst Oswaldo Cruz* 2006; 101: 239–243.
  73. Kordossis T, Avlami A, Velegriki A et al. First report of *Cryptococcus laurentii* meningitis and a fatal case of *Cryptococcus albidos* cryptococcaemia in AIDS patients. *Med Mycol* 1998; 36: 335–339.
  74. Garcia-Martos P, Noval JF, Garcia-Tapia A, Marin P, Puerto JL, Sepulveda A. Susceptibility to antifungal agents of *Cryptococcus* species of clinical interest. *Med Clin (Barc)* 2002; 119: 211–213.
  75. Kassamali H, Anaissie E, Ro J et al. Disseminated *Geotrichum candidum* infection. *J Clin Microbiol* 1987; 25: 1782–1783.
  76. Anaissie E, Bodey GP, Kantarjian H et al. New spectrum of fungal infections in patients with cancer. *Rev Infect Dis* 1989; 11: 369–378.
  77. Henrich TJ, Marty FM, Milner DA Jr, Thorner AR. Disseminated *Geotrichum candidum* infection in a patient with relapsed acute myelogenous leukemia following allogeneic stem cell transplantation and review of the literature. *Transpl Infect Dis* 2009; 11: 458–462.
  78. Sfakianakis A, Krasagakis K, Stefanidou M et al. Invasive cutaneous infection with *Geotrichum candidum* – sequential treatment with amphotericin B and voriconazole. *Med Mycol* 2007; 45: 81–84.
  79. Andre N, Coze C, Gentet JC, Perez R, Bernard JL. *Geotrichum candidum* septicemia in a child with hepatoblastoma. *Pediatr Infect Dis J* 2004; 23: 86.
  80. Ng KP, Soo-Hoo TS, Koh MT, Kwan PW. Disseminated *Geotrichum* infection. *Med J Malaysia* 1994; 49: 424–426.
  81. Wildfeuer A, Seidl HP, Paule I, Haberreiter A. *In vitro* evaluation of voriconazole against clinical isolates of yeasts, moulds and dermatophytes in comparison with itraconazole, ketoconazole, amphotericin B and griseofulvin. *Mycoses* 1998; 41: 309–319.
  82. Cordoba S, Vivot W, Bosco-Borgeat ME et al. Species distribution and susceptibility profile of yeasts isolated from blood cultures: results of a multicenter active laboratory-based surveillance study in Argentina. *Rev Argent Microbiol* 2011; 43: 176–185.
  83. Taj-Aldeen SJ, Doiphode SH, Han XY. *Kodamaea (Pichia) ohmeri* fungaemia in a premature neonate. *J Med Microbiol* 2006; 55: 237–239.
  84. Chakrabarti A, Rudramurthy SM, Kale P et al. *Kodamaea ohmeri* fungemia is not an uncommon entity in india: need for a systematic study. IDSA annual meeting 2010; M-652.
  85. De Barros JD, Do Nascimento SM, De Araujo FJ et al. *Kodamaea (Pichia) ohmeri* fungemia in a pediatric patient admitted in a public hospital. *Med Mycol* 2009; 47: 775–779.
  86. Al-Sweih N, Khan ZU, Ahmad S et al. *Kodamaea ohmeri* as an emerging pathogen: a case report and review of the literature. *Med Mycol* 2011; 49: 766–770.
  87. Sundaram PS, Bijulal S, Tharakan JA, Antony M. *Kodamaea ohmeri* tricuspid valve endocarditis with right ventricular inflow obstruction in a neonate with structurally normal heart. *Ann Pediatr Cardiol* 2011; 4: 77–80.
  88. Santino I, Bono S, Borruso L et al. *Kodamaea ohmeri* isolate from two immunocompromised patients: first report in Italy. *Mycoses* 2013; 56: 179–181.
  89. Menon T, Herrera M, Periasamy S, Palanivelu V, Sikhamani R, Wickes B. Oral candidiasis caused by *Kodamaea ohmeri* in a HIV patient in Chennai, India. *Mycoses* 2010; 53: 458–459.
  90. Mahfouz RA, Otrrock ZK, Mehawej H, Farhat F. *Kodamaea (Pichia) ohmeri* fungemia complicating acute myeloid leukaemia in a patient with haemochromatosis. *Pathology* 2008; 40: 99–101.
  91. Chiu CH, Wang YC, Shang ST, Chang FY. *Kodamaea ohmeri* fungaemia successfully treated with caspofungin. *Int J Antimicrob Agents* 2010; 35: 98–99.
  92. Yang BH, Peng MY, Hou SJ, Sun JR, Lee SY, Lu JJ. Fluconazole-resistant *Kodamaea ohmeri* fungemia associated with cellulitis: case report and review of the literature. *Int J Infect Dis* 2009; 13: e493–e497.
  93. Shaaban H, Choo HF, Boghossian J, Perez G. *Kodamaea ohmeri* fungemia in an immunocompetent patient treated with micafungin: case report and review of the literature. *Mycopathologia* 2010; 170: 223–228.
  94. Lee JS, Shin JH, Kim MN et al. *Kodamaea ohmeri* isolates from patients in a university hospital: identification, antifungal susceptibility, and pulsed-field gel electrophoresis analysis. *J Clin Microbiol* 2007; 45: 1005–1010.
  95. Otag F, Kuyucu N, Erturan Z, Sen S, Emekdas G, Sugita T. An outbreak of *Pichia ohmeri* infection in the paediatric intensive care unit: case reports and review of the literature. *Mycoses* 2005; 48: 265–269.
  96. Reina JP, Larone DH, Sabetta JR, Krieger KK, Hartman BJ. *Pichia ohmeri* prosthetic valve endocarditis and review of the literature. *Scand J Infect Dis* 2002; 34: 140–141.
  97. Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegriki A. The *Malassezia* genus in skin and systemic diseases. *Clin Microbiol Rev* 2012; 25: 106–141.
  98. Chang HJ, Miller HL, Watkins N et al. An epidemic of *Malassezia pachydermatis* in an intensive care nursery associated with colonization of health care workers' pet dogs. *N Engl J Med* 1998; 338: 706–711.
  99. Gueho E, Simmons RB, Pruitt WR, Meyer SA, Ahearn DG. Association of *Malassezia pachydermatis* with systemic infections of humans. *J Clin Microbiol* 1987; 25: 1789–1790.
  100. Morris DO. *Malassezia pachydermatis* carriage in dog owners. *Emerg Infect Dis* 2005; 11: 83–88.
  101. Chryssanthou E, Broberger U, Petrini B. *Malassezia pachydermatis* fungaemia in a neonatal intensive care unit. *Acta Paediatr* 2001; 90: 323–327.
  102. Mickelsen PA, Viano-Paulson MC, Stevens DA, Diaz PS. Clinical and microbiological features of infection with *Malassezia pachydermatis* in high-risk infants. *J Infect Dis* 1988; 157: 1163–1168.
  103. van Belkum A, Boekhout T, Bosboom R. Monitoring spread of *Malassezia* infections in a neonatal intensive care unit by PCR-mediated genetic typing. *J Clin Microbiol* 1994; 32: 2528–2532.
  104. Redline RW, Redline SS, Boxerbaum B, Dahms BB. Systemic *Malassezia furfur* infections in patients receiving intralipid therapy. *Hum Pathol* 1985; 16: 815–822.
  105. Morrison VA, Weisdorf DJ. The spectrum of *Malassezia* infections in the bone marrow transplant population. *Bone Marrow Transplant* 2000; 26: 645–648.
  106. Tragiannidis A, Bisping G, Koehler G, Groll AH. Minireview: *Malassezia* infections in immunocompromised patients. *Mycoses* 2010; 53: 187–195.
  107. Marcon MJ, Powell DA, Durrell DE. Methods for optimal recovery of *Malassezia furfur* from blood culture. *J Clin Microbiol* 1986; 24: 696–700.
  108. Nelson SC, Yau YC, Richardson SE, Matlow AG. Improved detection of *Malassezia* species in lipid-supplemented Peds Plus blood culture bottles. *J Clin Microbiol* 1995; 33: 1005–1007.
  109. Guillot J, Gueho E. The diversity of *Malassezia* yeasts confirmed by rRNA sequence and nuclear DNA comparisons. *Antonie Van Leeuwenhoek* 1995; 67: 297–314.
  110. Sugita T, Suzuki M, Goto S et al. Quantitative analysis of the cutaneous *Malassezia* microbiota in 770 healthy Japanese by age and gender using a real-time PCR assay. *Med Mycol* 2010; 48: 229–233.
  111. Gupta AK, Boekhout T, Theelen B, Summerbell R, Batra R. Identification and typing of *Malassezia* species by amplified fragment length polymorphism and sequence analyses of the internal transcribed spacer and large-subunit regions of ribosomal DNA. *J Clin Microbiol* 2004; 42: 4253–4260.
  112. Gueho-Kellermann E, Boekhout T, Begerow D. Biodiversity, phylogeny and ultrastructure. In: Boekhout T, Gueho E, Mayser P, Velegriki

- A, eds. *Malassezia* and the skin. Science and clinical practice. Heidelberg: Springer, 2010; 17–63.
113. Cafarchia C, Figueredo LA, Favuzzi V et al. Assessment of the antifungal susceptibility of *Malassezia pachydermatis* in various media using a CLSI protocol. *Vet Microbiol* 2012; 159: 536–540.
114. Jesus FPK, Lautert C, Zanette RA et al. *In vitro* susceptibility of fluconazole-susceptible and -resistant isolates of *Malassezia pachydermatis* against azoles. *Vet Microbiol* 2011; 152: 161–164.
115. Cafarchia C, Figueredo LA, Iatta R, Colao V, Montagna MT, Otranto D. *In vitro* evaluation of *Malassezia pachydermatis* susceptibility to azole compounds using E-test and CLSI microdilution methods. *Med Mycol* 2012; 50: 795–801.
116. Ashbee HR. Update on the genus *Malassezia*. *Med Mycol* 2007; 45: 287–303.
117. Devlin RK. Invasive fungal infections caused by *Candida* and *Malassezia* species in the neonatal intensive care unit. *Adv Neonatal Care* 2006; 6: 68–77.
118. Gerin M, Mahloui N, Elie C et al. Therapeutic drug monitoring of voriconazole after intravenous administration in infants and children with primary immunodeficiency. *Ther Drug Monit* 2011; 33: 464–466.
119. Pieper S, Kolve H, Gumbinger HG, Goletz G, Würthwein G, Groll AH. Monitoring of voriconazole plasma concentrations in immunocompromised paediatric patients. *J Antimicrob Chemother* 2012; 67: 2717–2724.
120. Bartelink IH, Wolfs T, Jonker M et al. Highly variable plasma concentrations of voriconazole in pediatric hematopoietic stem cell transplantation patients. *Antimicrob Agents Chemother* 2013; 57: 235–240.
121. Sugita T, Takashima M, Poonwan N et al. The first isolation of ustilaginomycetous anamorphic yeasts, *Pseudozyma* species, from patients' blood and a description of two new species: *P. parantarctica* and *P. thailandica*. *Microbiol Immunol* 2003; 47: 183–190.
122. Lin SS, Pranikoff T, Smith SF et al. Central venous catheter infection associated with *Pseudozyma aphidis* in a child with short gut syndrome. *J Med Microbiol* 2008; 57: 516–518.
123. Prakash A, Wankhede S, Singh PK et al. First neonatal case of fungemia due to *Pseudozyma aphidis* and a global literature review. *Mycoses* 2013. doi: 10.1111/myc.12098. [Epub ahead of print].
124. de Carvalho Parahym AMR, da Silva CM, Domingos IDF et al. Pulmonary infection due to *Pseudozyma aphidis* in a patient with Burkitt lymphoma: first case report. *Diagn Microbiol Infect Dis* 2013; 75: 104–106.
125. Hwang S, Kim J, Yoon S et al. First report of brain abscess associated with *Pseudozyma* species in a patient with astrocytoma. *Korean J Lab Med* 2013; 30: 284–288.
126. Chen B, Zhu LY, Xuan X et al. Isolation of both *Pseudozyma aphidis* and *Nocardia otitidiscaviarum* from a mycetoma on the leg. *Int J Dermatol* 2011; 50: 714–719.
127. Wirth F, Goldani LZ. Epidemiology of *Rhodotorula*: an emerging pathogen. *Interdiscip Perspect Infect Dis* 2012; 2012: 465717.
128. Sampaio JP. *Rhodotorula* Harrison (1928). In: Kurtzman CP, Fell JW, Boekhout T, eds. *The yeasts, a taxonomic study*, 5th edn. Amsterdam: Elsevier, 2011; 1873–1927.
129. Biswas SK, Yokoyama K, Nishimura K, Miyaji M. Molecular phylogenetics of the genus *Rhodotorula* and related basidiomycetous yeasts inferred from the mitochondrial cytochrome b gene. *Int J Syst Evol Microbiol* 2001; 51: 1191–1199.
130. Tuon FF, Costa SF. *Rhodotorula* infection. A systematic review of 128 cases from literature. *Rev Iberoam Micol* 2008; 25: 135–140.
131. Garcia-Suarez J, Gomez-Herruz P, Cuadros JA, Burgaleta C. Epidemiology and outcome of *Rhodotorula* infection in haematological patients. *Mycoses* 2011; 54: 318–324.
132. Tuon FF, de Almeida GM, Costa SF. Central venous catheter-associated fungemia due to *Rhodotorula* spp. – a systematic review. *Med Mycol* 2007; 45: 441–447.
133. Miceli MH, Diaz JA, Lee SA. Emerging opportunistic yeast infections. *Lancet Infect Dis* 2011; 11: 142–151.
134. Lunardi LW, Aquino VR, Zimerman RA, Goldani LZ. Epidemiology and outcome of *Rhodotorula* fungemia in a tertiary care hospital. *Clin Infect Dis* 2006; 43: e60–e63.
135. Fores R, Ramos A, Orden B et al. *Rhodotorula* species fungaemia causes low mortality in haematopoietic stem-cell transplantation. A case report and review. *Mycoses* 2012; 55: e158–e162.
136. Perniola R, Faneschi ML, Manso E et al. *Rhodotorula mucilaginosa* outbreak in neonatal intensive care unit: microbiological features, clinical presentation, and analysis of related variables. *Eur J Clin Microbiol Infect Dis* 2006; 25: 193–196.
137. Gyaurgieva OH, Bogomolova TS, Gorshkova GI. Meningitis caused by *Rhodotorula rubra* in an HIV-infected patient. *J Med Vet Mycol* 1996; 34: 357–359.
138. Unal A, Koc AN, Sipahioglu MH et al. CAPD-related peritonitis caused by *Rhodotorula mucilaginosa*. *Perit Dial Int* 2009; 29: 581–582.
139. Maeder M, Vogt PR, Schaer G, von Graevenitz A, Gunthard HF. Aortic homograft endocarditis caused by *Rhodotorula mucilaginosa*. *Infection* 2003; 31: 181–183.
140. Baradkar VP, Kumar S. Meningitis caused by *Rhodotorula mucilaginosa* in human immunodeficiency virus seropositive patient. *Ann Indian Acad Neurol* 2008; 11: 245–247.
141. Elias ML, Soliman AK, Mahoney FJ et al. Isolation of *Cryptococcus*, *Candida*, *Aspergillus*, *Rhodotorula* and *Nocardia* from meningitis patients in Egypt. *J Egypt Public Health Assoc* 2009; 84: 169–181.
142. Savini V, Sozio F, Catavittello C et al. Femoral prosthesis infection by *Rhodotorula mucilaginosa*. *J Clin Microbiol* 2008; 46: 3544–3545.
143. Shinde RS, Mantur BG, Patil G, Parande MV, Parande AM. Meningitis due to *Rhodotorula glutinis* in an HIV infected patient. *Indian J Med Microbiol* 2008; 26: 375–377.
144. Villar JM, Velasco CG, Delgado JD. Fungemia due to *Rhodotorula mucilaginosa* in an immunocompetent, critically ill patient. *J Infect Chemother* 2012; 18: 581–583.
145. Pinna A, Carta F, Zanetti S, Sanna S, Sechi LA. Endogenous *Rhodotorula minuta* and *Candida albicans* endophthalmitis in an injecting drug user. *Br J Ophthalmol* 2001; 85: 754.
146. Spiliopoulou A, Anastassiou ED, Christofidou M. *Rhodotorula* fungemia of an intensive care unit patient and review of published cases. *Mycopathologia* 2012; 174: 301–309.
147. Mori T, Nakamura Y, Kato J et al. Fungemia due to *Rhodotorula mucilaginosa* after allogeneic hematopoietic stem cell transplantation. *Transpl Infect Dis* 2012; 14: 91–94.
148. Garcia-Suarez J, Gomez-Herruz P, Cuadros JA, Guillen H, Burgaleta C. *Rhodotorula mucilaginosa* catheter-related fungaemia in a patient with multiple myeloma. *Mycoses* 2011; 54: e214–e216.
149. Poikonen E, Lyytikäinen O, Anttila VJ et al. Secular trend in candidemia and the use of fluconazole in Finland, 2004–2007. *BMC Infect Dis* 2010; 10: 312.
150. Khodavaisy S, Nabili M, Davari B, Vahedi M. Evaluation of bacterial and fungal contamination in the health care workers' hands and rings in the intensive care unit. *J Prev Med Hyg* 2011; 52: 215–218.
151. Pfaller MA, Diekema DJ. Rare and emerging opportunistic fungal pathogens: concern for resistance beyond *Candida albicans* and *Aspergillus fumigatus*. *J Clin Microbiol* 2004; 42: 4419–4431.
152. Hall L, Le Febvre KM, Deml SM, Wohlfel SL, Wengenack NL. Evaluation of the yeast traffic light PNA FISH probes for identification of *Candida* species from positive blood cultures. *J Clin Microbiol* 2012; 50: 1446–1448.
153. Diekema DJ, Petroelje B, Messer SA, Hollis RJ, Pfaller MA. Activities of available and investigational antifungal agents against *Rhodotorula* species. *J Clin Microbiol* 2005; 43: 476–478.
154. Nunes JM, Bizerra FC, Ferreira RC, Colombo AL. Molecular identification, antifungal susceptibility profile and biofilm formation of clinical and environmental *Rhodotorula* spp. isolates. *Antimicrob Agents Chemother* 2013; 57: 382–389.

155. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope WW; The European Committee on Antimicrobial Susceptibility Testing Subcommittee on Antifungal Susceptibility Testing (EUCAST-AFST). EUCAST technical note on *Aspergillus* and amphotericin-B, itraconazole, and posaconazole. *Clin Microbiol Infect* 2012; 17: E248–E250.
156. Subcommittee on Antifungal Susceptibility Testing (AFST) of the ESCMID European Committee for Antimicrobial Susceptibility Testing (EUCAST). EUCAST technical note on voriconazole. *Clin Microbiol Infect* 2008; 14: 985–987.
157. Hope WW, Cuenca-Estrella M, Lass-Flörl C, Arendrup MC. EUCAST technical note on voriconazole and *Aspergillus* spp. *Clin Microbiol Infect* 2013; 19: E278–E280.
158. Preney L, Theraud M, Guiguen C, Gangneux JP. Experimental evaluation of antifungal and antiseptic agents against *Rhodotorula* spp. *Mycoses* 2003; 46: 492–495.
159. Gomez-Lopez A, Mellado E, Rodriguez-Tudela JL, Cuenca-Estrella M. Susceptibility profile of 29 clinical isolates of *Rhodotorula* spp. and literature review. *J Antimicrob Chemother* 2005; 55: 312–316.
160. Caira M, Treccarichi EM, Tumbarello M, Leone G, Pagano L. Uncommon yeast infections in hematological patients: from diagnosis to treatment. *Expert Rev Anti Infect Ther* 2011; 9: 1067–1075.
161. Anatoliotaki M, Mantadakis E, Galanakis E, Samonis G. *Rhodotorula* species fungemia: a threat to the immunocompromised host. *Clin Lab* 2003; 49: 49–55.
162. de Zoysa JR, Searle M, Lynn KL, Robson RA. Successful treatment of CAPD peritonitis caused by *Rhodotorula mucilaginosa*. *Perit Dial Int* 2001; 21: 627–628.
163. Asim M, Meigh JA, Farr MJ. Fungal peritonitis caused by *Candida parapsilosis* and subsequently *Rhodotorula glutinis* in a CAPD patient. *Perit Dial Int* 1999; 19: 183.
164. Samonis G, Anatoliotaki M, Apostolou H, Maraki S, Mavroudis D, Georgoulis V. Transient fungemia due to *Rhodotorula rubra* in a cancer patient: case report and review of the literature. *Infection* 2001; 29: 173–176.
165. Edwards-Ingram LC, Gent ME, Hoyle DC, Hayes A, Stateva LI, Oliver SG. Comparative genomic hybridization provides new insights into the molecular taxonomy of the *Saccharomyces sensu stricto* complex. *Genome Res* 2004; 14: 1043–1051.
166. Edwards-Ingram L, Gitsham P, Burton N et al. Genotypic and physiological characterization of *Saccharomyces boulardii*, the probiotic strain of *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 2007; 73: 2458–2467.
167. Riaz M, Alam S, Malik A, Ali SM. Efficacy and safety of *Saccharomyces boulardii* in acute childhood diarrhea: a double blind randomised controlled trial. *Indian J Pediatr* 2012; 79: 478–482.
168. de Llanos R, Llopis S, Molero G, Querol A, Gil C, Fernandez-Espinar MT. *In vivo* virulence of commercial *Saccharomyces cerevisiae* strains with pathogenicity-associated phenotypic traits. *Int J Food Microbiol* 2011; 144: 393–399.
169. Czeruckat D, Piche P, Rampal P. Review article: yeast as probiotics – *Saccharomyces boulardii*. *Aliment Pharmacol Ther* 2007; 26: 767–778.
170. Richter SS, Galask RP, Messer SA, Hollis RJ, Diekema DJ, Pfaller MA. Antifungal susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent cases. *J Clin Microbiol* 2005; 43: 2155–2162.
171. Thygesen JB, Glerup H, Tarp B. *Saccharomyces boulardii* fungemia caused by treatment with a probiotic. *BMJ Case Rep* 2012. doi: 10.1136/bcr.2011.4412
172. Stefanatou E, Kompoti M, Paridou A et al. Probiotic sepsis due to *Saccharomyces* fungaemia in a critically ill burn patient. *Mycoses* 2011; 54: e643–e646.
173. Bassetti S, Frei R, Zimmerli W. Fungemia with *Saccharomyces cerevisiae* after treatment with *Saccharomyces boulardii*. *Am J Med* 1998; 105: 71–72.
174. de Llanos R, Querol A, Peman J, Gobernado M, Fernandez-Espinar MT. Food and probiotic strains from the *Saccharomyces cerevisiae* species as a possible origin of human systemic infections. *Int J Food Microbiol* 2006; 110: 286–290.
175. Herbrecht R, Nivoix Y. *Saccharomyces cerevisiae* fungemia: an adverse effect of *Saccharomyces boulardii* probiotic administration. *Clin Infect Dis* 2005; 40: 1635–1637.
176. Munoz P, Bouza E, Cuenca-Estrella M et al. *Saccharomyces cerevisiae* fungemia: an emerging infectious disease. *Clin Infect Dis* 2005; 40: 1625–1634.
177. Graf C, Gavazzi G. *Saccharomyces cerevisiae* fungemia in an immunocompromised patient not treated with *Saccharomyces boulardii* preparation. *J Infect* 2007; 54: 310–311.
178. Enache-Angoulvant A, Hennequin C. Invasive *Saccharomyces* infection: a comprehensive review. *Clin Infect Dis* 2005; 41: 1559–1568.
179. Rimek D, Redetzke K, Singh J, Heinrich K, Kappe R. Performance des Candida-Mannan-Antigennachweises bei Patienten mit Fungämien. *Mycoses* 2004; 47: 23–26.
180. Yoshida M, Obayashi T, Iwama A et al. Detection of plasma (1→3)- $\beta$ -D-glucan in patients with *Fusarium*, *Trichosporon*, *Saccharomyces* and *Acremonium* fungaemias. *J Med Vet Mycol* 1997; 35: 371–374.
181. Thompson GR III, Wiederhold NP, Sutton DA, Fothergill A, Patterson TF. *In vitro* activity of isavuconazole against *Trichosporon*, *Rhodotorula*, *Geotrichum*, *Saccharomyces* and *Pichia* species. *J Antimicrob Chemother* 2009; 64: 79–83.
182. Lolis N, Veldekis D, Moraitou H et al. *Saccharomyces boulardii* fungaemia in an intensive care unit patient treated with caspofungin. *Crit Care* 2008; 12: 414.
183. Choi G, Meijer SL, Hazenberg MD. Disseminated bread yeast fungaemia in a baker's wife with acute myeloid leukaemia. *Br J Haematol* 2012; 158: 298.
184. Tiballi RN, Spiegel JE, Zarins LT, Kauffman CA. *Saccharomyces cerevisiae* infections and antifungal susceptibility studies by colorimetric and broth macrodilution methods. *Diagn Microbiol Infect Dis* 1995; 23: 135–140.
185. Hamoud S, Keidar Z, Hayek T. Recurrent *Saccharomyces cerevisiae* fungemia in an otherwise healthy patient. *Isr Med Assoc J* 2011; 13: 575–576.
186. Gadea I, Cuenca-Estrella M, Prieto E et al. Genotyping and antifungal susceptibility profile of *Dipodascus capitatus* isolates causing disseminated infection in seven hematological patients of a tertiary hospital. *J Clin Microbiol* 2004; 42: 1832–1836.
187. Aalbaek B, Stenderup J, Jensen HE, Valbak J, Nylin B, Huda A. Mycotic and algal bovine mastitis in Denmark. *APMIS* 1994; 102: 451–456.
188. Döğen A, Kaplan E, Öksüz Z, Serin MS, İkit M, de Hoog GS. Dishwashers are a major source of human opportunistic yeast-like fungi in indoor environments in Mersin, Turkey. *Med Mycol* 2013; 51: 493–498.
189. Martino P, Girmenia C, Venditti M et al. Spontaneous pneumothorax complicating pulmonary mycetoma in patients with acute leukemia. *Rev Infect Dis* 1990; 12: 611–617.
190. Birrenbach T, Bertschy S, Aebersold F et al. Emergence of *Blastoschizomyces capitatus* yeast infections, Central Europe. *Emerg Infect Dis* 2012; 18: 98–101.
191. Christakis G, Perlorentzou S, Aslanidou M, Megalaki A, Velegraki A. Fatal *Blastoschizomyces capitatus* sepsis in a neutropenic patient with acute myeloid leukemia: first documented case from Greece. *Mycoses* 2005; 48: 216–220.
192. Girmenia C, Pagano L, Martino B et al. Invasive infections caused by *Trichosporon* species and *Geotrichum capitatum* in patients with hematological malignancies: a retrospective multicenter study from Italy and review of the literature. *J Clin Microbiol* 2005; 43: 1818–1828.
193. Martino R, Salavert M, Parody R et al. *Blastoschizomyces capitatus* infection in patients with leukemia: report of 26 cases. *Clin Infect Dis* 2004; 38: 335–341.
194. DeMaio J, Colman L. The use of adjuvant interferon- $\gamma$  therapy for hepatosplenic *Blastoschizomyces capitatus* infection in a patient with leukemia. *Clin Infect Dis* 2000; 31: 822–824.

195. Pagano L, Morace G, Ortu-La BE, Sanguinetti M, Leone G. Adjuvant therapy with rhGM-CSF for the treatment of *Blastoschizomyces capitatus* systemic infection in a patient with acute myeloid leukemia. *Ann Hematol* 1996; 73: 33–34.
196. Cofrancesco E, Viviani MA, Boschetti C, Tortorano AM, Balzani A, Castagnone D. Treatment of chronic disseminated *Geotrichum capitatum* infection with high cumulative dose of colloidal amphotericin B and itraconazole in a leukaemia patient. *Mycoses* 1995; 38: 377–384.
197. Oravcova E, Lacka J, Drgona L et al. Funguria in cancer patients: analysis of risk factors, clinical presentation and outcome in 50 patients. *Infection* 1996; 24: 319–323.
198. Schiemann R, Glasmacher A, Bailly E et al. *Geotrichum capitatum* septicaemia in neutropenic patients: case report and review of the literature. *Mycoses* 1998; 41: 113–116.
199. Cheung MY, Chiu NC, Chen SH, Liu HC, Ou CT, Liang DC. Mandibular osteomyelitis caused by *Blastoschizomyces capitatus* in a child with acute myelogenous leukemia. *J Formos Med Assoc* 1999; 98: 787–789.
200. Celik A, Ozaras R, Kantarcioglu S, Mert A, Tabak F, Ozturk R. Spondylodiscitis due to an emergent fungal pathogen: *Blastoschizomyces capitatus*, a case report and review of the literature. *Rheumatol Int* 2009; 29: 1237–1241.
201. Lapusan S, Dimicoli S, Marjanovic Z et al. Immune reconstitution inflammatory syndrome following allo-SCT in a patient with *Dipodascus capitatus* spondylodiscitis. *Bone Marrow Transplant* 2011; 46: 1265–1267.
202. D'Antonio D, Mazzoni A, Iacone A et al. Emergence of fluconazole-resistant strains of *Blastoschizomyces capitatus* causing nosocomial infections in cancer patients. *J Clin Microbiol* 1996; 34: 753–755.
203. Gurgui M, Sanchez F, March F et al. Nosocomial outbreak of *Blastoschizomyces capitatus* associated with contaminated milk in a haematological unit. *J Hosp Infect* 2011; 78: 274–278.
204. Polachek I, Salkin IF, Kitzes-Cohen R, Raz R. Endocarditis caused by *Blastoschizomyces capitatus* and taxonomic review of the genus. *J Clin Microbiol* 1992; 30: 2318–2322.
205. Wills TS, Degryse A, Lavina J, Sinnott JT. *Blastoschizomyces capitatus* pneumonia in an immunocompetent male. *South Med J* 2004; 97: 702–704.
206. Mathews MS, Sen S. *Blastoschizomyces capitatus* infection after contamination of fluids for intravenous application. *Mycoses* 1998; 41: 427–428.
207. Naficy AB, Murray HW. Isolated meningitis caused by *Blastoschizomyces capitatus*. *J Infect Dis* 1990; 161: 1041–1042.
208. Giacchino M, Chiappello N, Bezzio S et al. *Aspergillus* galactomannan enzyme-linked immunosorbent assay cross-reactivity caused by invasive *Geotrichum capitatum*. *J Clin Microbiol* 2006; 44: 3432–3434.
209. Bonini A, Capatti C, Parmeggiani M et al. Galactomannan detection in *Geotrichum capitatum* invasive infections: report of 2 new cases and review of diagnostic options. *Diagn Microbiol Infect Dis* 2008; 62: 450–452.
210. Girmenia C, Pizzarelli G, D'Antonio D, Cristini F, Martino P. *In vitro* susceptibility testing of *Geotrichum capitatum*: comparison of the E-test, disk diffusion, and Sensititre colorimetric methods with the NCCLS M27-A2 broth microdilution reference method. *Antimicrob Agents Chemother* 2003; 47: 3985–3988.
211. Chittick P, Palavecino EL, Delashmitt B, Evans J, Peacock JE. Case of fatal *Blastoschizomyces capitatus* infection occurring in a patient receiving empiric micafungin therapy. *Antimicrob Agents Chemother* 2009; 53: 5306–5307.
212. Schuermans C, van Bergen M, Coorevits L et al. Breakthrough *Saprochaete capitata* infections in patients receiving echinocandins: case report and review of the literature. *Med Mycol* 2011; 49: 414–418.
213. Serena C, Marine M, Marimon R, Pastor FJ, Guarro J. Effect of antifungal treatment in a murine model of blastoschizomycosis. *Int J Antimicrob Agents* 2007; 29: 79–83.
214. Ikuta K, Torimoto Y, Yamamoto M et al. Successful treatment of systemic *Geotrichum capitatum* infection by liposomal amphotericin-B, itraconazole, and voriconazole in a Japanese man. *Intern Med* 2010; 49: 2499–2503.
215. Etienne A, Datry A, Gaspar N, Morel V, Delabesse E, Lmimouni B, et al. Successful treatment of disseminated *Geotrichum capitatum* infection with a combination of caspofungin and voriconazole in an immunocompromised patient. *Mycoses* 2008; 51: 270–272.
216. Fianchi L, Montini L, Caira M et al. Combined voriconazole plus caspofungin therapy for the treatment of probable *Geotrichum* pneumonia in a leukemia patient. *Infection* 2008; 36: 65–67.
217. Perez-Sanchez I, Anguita J, Rabadan PM et al. *Blastoschizomyces capitatus* infection in acute leukemia patients. *Leuk Lymphoma* 2000; 39: 209–212.
218. Hamamoto M, Boekhout T, Nakase T. *Sporobolomyces Klyver & van Niel* (1914). In: Kurtzman CP, Fell JW, Boekhout T, eds. *The yeasts, a taxonomic study*, 5th edn. Amsterdam: Elsevier, 2011; 1929–1990.
219. Fairs A, Wardlaw AJ, Thompson J, Pashley CH. Guidelines on ambient intramural airborne fungal spores. *J Investig Allergol Clin Immunol* 2010; 20: 490–498.
220. Moore JE, Xu J, Millar BC, Elshibly S. Edible dates (*Phoenix dactylifera*), a potential source of *Cladosporium cladosporioides* and *Sporobolomyces roseus*: implications for public health. *Mycopathologia* 2002; 154: 25–28.
221. Watanabe M, Tsutsumi F, Konuma R et al. Quantitative analysis of mycoflora on commercial domestic fruits in Japan. *J Food Prot* 2011; 74: 1488–1499.
222. Morris JT, Beckius M, McAllister CK. *Sporobolomyces* infection in an AIDS patient. *J Infect Dis* 1991; 164: 623–624.
223. Plazas J, Portilla J, Boix V, Perez-Mateo M. *Sporobolomyces salmonicolor* lymphadenitis in an AIDS patient. Pathogen or passenger? *AIDS* 1994; 8: 387–388.
224. McNicholas S, McDermott H, Power L et al. *Sporobolomyces roseus* in the cerebrospinal fluid of an immunocompetent patient – to treat or not to treat? *J Med Microbiol* 2012; 61: 295–296.
225. Sharma V, Shankar J, Kotamarthi V. Endogenous endophthalmitis caused by *Sporobolomyces salmonicolor*. *Eye* 2005; 20: 945–946.
226. Bergman AG, Kauffman CA. Dermatitis due to *Sporobolomyces* infection. *Arch Dermatol* 1984; 120: 1059–1060.
227. Cockcroft DW, Berscheid BA, Ramshaw IA, Dolovich J. *Sporobolomyces*: a possible cause of extrinsic allergic alveolitis. *J Allergy Clin Immunol* 1983; 72: 305–309.
228. Seuri M, Husman K, Kinnunen H et al. An outbreak of respiratory diseases among workers at a water-damaged building – a case report. *Indoor Air* 2000; 10: 138–145.
229. Bross JE, Manning P, Kacian D, Talbot GH. Pseudomeningitis caused by *Sporobolomyces salmonicolor*. *Am J Infect Control* 1986; 14: 220–223.
230. Espinel-Ingroff A. *In vitro* activity of the new triazole voriconazole (UK-109,496) against opportunistic filamentous and dimorphic fungi and common and emerging yeast pathogens. *J Clin Microbiol* 1998; 36: 198–202.
231. Colombo AL, Padovan AC, Chaves GM. Current knowledge of *Trichosporon* spp. and Trichosporonosis. *Clin Microbiol Rev* 2011; 24: 682–700.
232. Ruan SY, Chien JY, Hsueh PR. Invasive trichosporonosis caused by *Trichosporon asahii* and other unusual *Trichosporon* species at a medical center in Taiwan. *Clin Infect Dis* 2009; 49: e11–e17.
233. Fournier S, Pavageau W, Feuillhade M et al. Use of voriconazole to successfully treat disseminated *Trichosporon asahii* infection in a patient with acute myeloid leukaemia. *Eur J Clin Microbiol Infect Dis* 2002; 21: 892–896.
234. Wynne SM, Kwon-Chung KJ, Shea YR et al. Invasive infection with *Trichosporon inkin* in 2 siblings with chronic granulomatous disease. *J Allergy Clin Immunol* 2004; 114: 1418–1424.
235. Hickey PV, Sutton DA, Fothergill AW et al. *Trichosporon mycotoxivorans*, a novel respiratory pathogen in patients with cystic fibrosis. *J Clin Microbiol* 2009; 47: 3091–3097.

236. Kontoyiannis DP, Torres HA, Chagua M et al. Trichosporonosis in a tertiary care cancer center: risk factors, changing spectrum and determinants of outcome. *Scand J Infect Dis* 2004; 36: 564–569.
237. Matsue K, Uryu H, Koseki M, Asada N, Takeuchi M. Breakthrough trichosporonosis in patients with hematologic malignancies receiving micafungin. *Clin Infect Dis* 2006; 42: 753–757.
238. Suzuki K, Nakase K, Kyo T et al. Fatal *Trichosporon* fungemia in patients with hematologic malignancies. *Eur J Haematol* 2010; 84: 441–447.
239. Bayramoglu G, Sonmez M, Tosun I, Aydin K, Aydin F. Breakthrough *Trichosporon asahii* fungemia in neutropenic patient with acute leukemia while receiving caspofungin. *Infection* 2008; 36: 68–70.
240. Pagano L, Caira M, Candoni A et al. The epidemiology of fungal infections in patients with hematologic malignancies: the SEIFEM-2004 study. *Haematologica* 2006; 91: 1068–1075.
241. Walsh TJ, Groll A, Hiemenz J, Fleming R, Roilides E, Anaissie E. Infections due to emerging and uncommon medically important fungal pathogens. *Clin Microbiol Infect* 2004; 10(Suppl 1): 48–66.
242. Panagopoulou P, Evdoridou J, Bibashi E et al. *Trichosporon asahii*: an unusual cause of invasive infection in neonates. *Pediatr Infect Dis J* 2002; 21: 169–170.
243. Vashishtha VM, Mittal A, Garg A. A fatal outbreak of *Trichosporon asahii* sepsis in a neonatal intensive care unit. *Indian Pediatr* 2012; 49: 745–747.
244. Moreno S, Buzon L, Sanchez-Sousa A. *Trichosporon capitatum* fungemia and intravenous drug abuse. *Rev Infect Dis* 1987; 9: 1202–1203.
245. Liao Y, Hartmann T, Ao JH, Yang RY. Serum glucuronoxylomannan may be more appropriate for the diagnosis and therapeutic monitoring of *Trichosporon* fungemia than serum  $\beta$ -D-glucan. *Int J Infect Dis* 2012; 16: e638.
246. McManus EJ, Bozdech MJ, Jones JM. Role of the latex agglutination test for cryptococcal antigen in diagnosing disseminated infections with *Trichosporon beigelii*. *J Infect Dis* 1985; 151: 1167–1169.
247. McManus EJ, Jones JM. Detection of a *Trichosporon beigelii* antigen cross-reactive with *Cryptococcus neoformans* capsular polysaccharide in serum from a patient with disseminated *Trichosporon* infection. *J Clin Microbiol* 1985; 21: 681–685.
248. Campbell CK, Payne AL, Teall AJ, Brownell A, Mackenzie DW. Cryptococcal latex antigen test positive in patient with *Trichosporon beigelii* infection. *Lancet* 1985; 2: 43–44.
249. Melcher GP, Reed KD, Rinaldi MG, Lee JW, Pizzo PA, Walsh TJ. Demonstration of a cell wall antigen cross-reacting with cryptococcal polysaccharide in experimental disseminated trichosporonosis. *J Clin Microbiol* 1991; 29: 192–196.
250. Nakase K, Suzuki K, Kyo T, Kohara T, Sugawara Y, Katayama N. Is elevation of the serum  $\beta$ -D-glucan level a paradoxical sign for *Trichosporon* fungemia in patients with hematologic disorders? *Int J Infect Dis* 2012; 16: e2–e4.
251. Kushima H, Ishii H, Komiya K, Tokimatsu I, Kadota JI. Prognostic significance of serum  $\beta$ -D-glucan levels in 78 patients with *Trichosporon* fungemia. *Int J Infect Dis* 2013; 17: e134–e135.
252. Shinozaki M, Okubo Y, Sasai D et al. Development of a peptide nucleic acid probe to *Trichosporon* species and identification of trichosporonosis by use of *in situ* hybridization in formalin-fixed and paraffin-embedded (FFPE) sections. *J Clin Microbiol* 2013; 51: 295–298.
253. Hosoki K, Iwamoto S, Kumamoto T, Azuma E, Komada Y. Early detection of breakthrough trichosporonosis by serum PCR in a cord blood transplant recipient being prophylactically treated with voriconazole. *J Pediatr Hematol Oncol* 2008; 30: 917–919.
254. Spiess B, Seifarth W, Hummel M et al. DNA microarray-based detection and identification of fungal pathogens in clinical samples from neutropenic patients. *J Clin Microbiol* 2007; 45: 3743–3753.
255. Nagai H, Yamakami Y, Hashimoto A, Tokimatsu I, Nasu M. PCR detection of DNA specific for *Trichosporon* species in serum of patients with disseminated trichosporonosis. *J Clin Microbiol* 1999; 37: 694–699.
256. Araujo RM, Alastruey-Izquierdo A, Gomez-Lopez A, Rodriguez-Tudela JL, Cuenca-Estrella M. Molecular identification and susceptibility testing of *Trichosporon* isolates from a Brazilian hospital. *Rev Iberoam Micol* 2008; 25: 221–225.
257. Rodriguez-Tudela JL, Diaz-Guerra TM, Mellado E et al. Susceptibility patterns and molecular identification of *Trichosporon* species. *Antimicrob Agents Chemother* 2005; 49: 4026–4034.
258. Sugita T, Nakajima M, Ikeda R, Matsushima T, Shinoda T. Sequence analysis of the ribosomal DNA intergenic spacer I regions of *Trichosporon* species. *J Clin Microbiol* 2002; 40: 1826–1830.
259. Kolecka A, Khayhan K, Groenewald M et al. MALDI-TOF MS identification of medically relevant species of arthroconidial yeasts. *J Clin Microbiol* 2013; 51: 2491–2500.
260. Asada N, Uryu H, Koseki M, Takeuchi M, Komatsu M, Matsue K. Successful treatment of breakthrough *Trichosporon asahii* fungemia with voriconazole in a patient with acute myeloid leukemia. *Clin Infect Dis* 2006; 43: e39–e41.
261. Serena C, Gilgado F, Marine M, Pastor FJ, Guarro J. Efficacy of voriconazole in a guinea pig model of invasive trichosporonosis. *Antimicrob Agents Chemother* 2006; 50: 2240–2243.
262. Rodriguez-Tudela JL, Gomez-Lopez A, Alastruey-Izquierdo A, Mellado E, Bernal-Martinez L, Cuenca-Estrella M. Genotype distribution of clinical isolates of *Trichosporon asahii* based on sequencing of intergenic spacer I. *Diagn Microbiol Infect Dis* 2007; 58: 435–440.
263. Chagas-Neto TC, Chaves GM, Melo AS, Colombo AL. Bloodstream infections due to *Trichosporon* spp.: species distribution, *Trichosporon asahii* genotypes determined on the basis of ribosomal DNA intergenic spacer I sequencing, and antifungal susceptibility testing. *J Clin Microbiol* 2009; 47: 1074–1081.
264. Taj-Aldeen SJ, Al-Ansari N, El Shafei S et al. Molecular identification and susceptibility of *Trichosporon* species isolated from clinical specimens in Qatar: isolation of *Trichosporon dohaense* Taj-Aldeen, Meis & Boekhout sp. nov. *J Clin Microbiol* 2009; 47: 1791–1799.
265. Hoy J, Hsu KC, Rolston K, Hopfer RL, Luna M, Bodey GP. *Trichosporon beigelii* infection: a review. *Rev Infect Dis* 1986; 8: 959–967.
266. Walsh TJ, Melcher GP, Rinaldi MG et al. *Trichosporon beigelii*, an emerging pathogen resistant to amphotericin B. *J Clin Microbiol* 1990; 28: 1616–1622.
267. Marty FM, Barouch DH, Coakley EP, Baden LR. Disseminated trichosporonosis caused by *Trichosporon loubieri*. *J Clin Microbiol* 2003; 41: 5317–5320.
268. Tokimatsu I, Kushima H, Hashinaga K et al. The prophylactic effectiveness of various antifungal agents against the progression of trichosporonosis fungemia to disseminated disease in a neutropenic mouse model. *Int J Antimicrob Agents* 2007; 29: 84–88.
269. Hosokawa K, Yamazaki H, Mochizuki K et al. Successful treatment of *Trichosporon* fungemia in a patient with refractory acute myeloid leukemia using voriconazole combined with liposomal amphotericin B. *Transpl Infect Dis* 2012; 14: 184–187.
270. Gabriel F, Noel T, Accoceberry I. Fatal invasive trichosporonosis due to *Trichosporon loubieri* in a patient with T-lymphoblastic lymphoma. *Med Mycol* 2010; 49: 306–310.
271. Antachopoulos C, Papakonstantinou E, Dotis J et al. Fungemia due to *Trichosporon asahii* in a neutropenic child refractory to amphotericin B: clearance with voriconazole. *J Pediatr Hematol Oncol* 2005; 27: 283–285.
272. Groll AH, Giri N, Petratis V et al. Comparative efficacy and distribution of lipid formulations of amphotericin B in experimental *Candida albicans* infection of the central nervous system. *J Infect Dis* 2000; 182: 274–282.
273. Hamill RJ, Sobel JD, El-Sadr W et al. Comparison of 2 doses of liposomal amphotericin B and conventional amphotericin B deoxycholate for treatment of AIDS-associated acute cryptococcal meningitis: a randomized, double-blind clinical trial of efficacy and safety. *Clin Infect Dis* 2010; 51: 225–232.